



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>6</sup> : C07H 17/08, A61K 31/70</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/21869 (43) International Publication Date: 6 May 1999 (06.05.99)</p>
<p>(21) International Application Number: PCT/JP98/04876 (22) International Filing Date: 28 October 1998 (28.10.98) (30) Priority Data: 9/296822 29 October 1997 (29.10.97) JP (71) Applicant (for all designated States except US): TAISHO PHARMACEUTICAL CO., LTD. [JP/JP]; 24-1, Takata 3-chome, Toshima-ku, Tokyo 171-8633 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): ASAKA, Toshifumi [JP/JP]; 2843-2, Nukata, Konosu-shi, Saitama 365-0059 (JP). KASHIMURA, Masato [JP/JP]; Raionzugaden Higashi-omiya-ichibankan 802, 702-12, Shimacho, Omiya-shi, Saitama 330-0006 (JP). MATSUURA, Akiko [JP/JP]; Byukuresutoru 101, 38-1, Higashitokorozawa 2-chome, Tokorozawa-shi, Saitama 359-0021 (JP). SUGIMOTO, Tomohiro [JP/JP]; Yoshinodokushinryo 216, 350-1, Yoshinocho 1-chome, Omiya-shi, Saitama 330-0031 (JP). TANIKAWA, Tetsuya [JP/JP]; Aruteru Fuchu 402, 12-5, Honmachi 1-chome, Fuchu-shi, Tokyo 183-0027 (JP). ISHII, Takaaki [JP/JP]; 6-7, Miyamoto 1-chome, Urawa-shi, Saitama 336-0916 (JP).</p>	<p>(74) Agents: ASAMURA, Kiyoshi et al.; New Ohtemachi Building, Room 331, 2-1, Ohtemachi 2-chome, Chiyoda-ku, Tokyo 100-0004 (JP). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> With international search report.</p>	
<p>(54) Title: ERYTHROMYCIN A, 11,12-CARBAMATE DERIVATIVES</p>		
<div style="text-align: center;"> <p>(I)</p> </div>		
<p>(57) Abstract</p> <p>An erythromycin A derivative represented by formula (I) wherein n is an integer of 1 to 7, R<sup>1</sup> is a group represented by the formula: -SO<sub>2</sub>N(-R<sup>7</sup>)-R<sup>8</sup> or N-(R<sup>10</sup>)SO<sub>2</sub>R<sup>9</sup>, R<sup>2</sup> is a hydrogen atom, an alkyl group or a cinnamyl group, R<sup>3</sup> is a group represented by the formula: -OCO-CH<sub>2</sub>-R<sup>11</sup>, -OCO-R<sup>11</sup>, -OCO-NH-R<sup>11</sup>, -O-R<sup>11</sup> or -OCO-O-R<sup>11</sup>, R<sup>4</sup> is a hydrogen atom, or R<sup>3</sup> and R<sup>4</sup> together form an oxo group, and R<sup>5</sup> and R<sup>6</sup> are each a hydrogen atom or an alkyl group, or a pharmaceutically acceptable salt thereof has a strong antibacterial activity against not only known erythromycin-sensitive bacteria but also erythromycin-resistant bacteria.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## DESCRIPTION

## ERYTHROMYCIN A, 11,12-CARBAMATE DERIVATIVES

## 5 TECHNICAL FIELD

The present invention relates to novel derivatives of antibiotic erythromycin A.

## BACKGROUND ART

10 Erythromycin A is an antibiotic clinically widely used as an agent for treating infectious diseases caused by Gram-positive bacteria, mycoplasmas, etc. However, erythromycin A is decomposed by the gastric acid due to instability to acids, and thereby has a  
15 drawback of no constancy of movement in the body. Hitherto many erythromycin A derivatives have been prepared for the purpose of the improvement of the biological or pharmacological properties. For example, it is reported that 6-O-methylethromycin A derivatives  
20 have an improved stability to acids and have a superior in vivo antibacterial activity in comparison with erythromycin A when administered orally (U.S. Patent No. 4331803). Recently, it is also reported that 11,12-cyclic carbamate derivatives are prepared from 6-O-  
25 methylethromycin A as a starting material with the aim of expansion of antibacterial spectrum as well as stability to acids (EP. patent No. 487411 and US. Patent No. 4742049). In addition, the antibacterial activities

of the ester derivatives at the 3-position are also reported by some of the present inventors (EP. patent No.619320).


An object of the present invention is to  
5 provide post-generational macrolide antibiotics having a strong antibacterial activity against not only known erythromycin-sensitive bacteria but also erythromycin-resistant bacteria which recently are showing a tendency to increase.

10

#### DISCLOSURE OF THE INVENTION


The present inventors have found that the compounds which can be produced by introducing a sulfonamide group onto the alkyl group attached to the  
15 nitrogen atom of the 11,12-cyclic carbamate of erythromycin A and converting 3-position of the erythromycin A have a strong antibacterial activity against not only erythromycin-sensitive bacteria but also erythromycin-resistant bacteria, and thus the  
20 present invention has been accomplished.

The present invention relates to an erythromycin A derivative represented by the formula:




wherein  $n$  is an integer of 1 to 7,

$R^1$  is a group represented by the formula:



- 5 wherein  $R^7$  is a hydrogen atom, an alkyl group having 1 to 6 carbon atoms, a phenyl group, a phenyl group substituted by a nitro group or an alkoxy group having 1 to 3 carbon atoms, a pyridyl group, a pyridyl group substituted by 1 or 2 members selected from the group
- 10 consisting of an alkyl group having 1 to 6 carbon atoms; a halogen atom; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group and an amino group substituted by an alkyl group having 1 to 6 carbon atoms, a quinolyl group, or a quinolyl group
- 15 substituted by 1 or 2 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms; a halogen atom; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group and an amino group substituted by an alkyl group having 1 to
- 20 6 carbon atoms,  $R^8$  is a hydrogen atom or an alkyl group having 1 to 6 carbon atoms, or a group represented by the formula:



wherein  $R^9$  is an alkyl group having 1 to 6 carbon atoms, a dibenzofuranyl group, a thienyl group, a thienyl group substituted by a group selected from the group

5 consisting of a pyridyl group; an isoxazolyl group; a pyrimidinyl group and a pyrimidinyl group substituted by an alkoxy group having 1 to 6 carbon atoms or an alkylthio group having 1 to 6 carbon atoms, an isoxazolyl group, an isoxazolyl group substituted by 1


10 or 2 alkyl groups having 1 to 6 carbon atoms, an imidazolyl group, an imidazolyl group substituted by 1 to 3 alkyl groups having 1 to 6 carbon atoms, a benzothienyl group, a benzothienyl group substituted by 1 to 5 members selected from the group consisting of an

15 alkyl group having 1 to 6 carbon atoms and a halogen atom, a thiazolyl group, a thiazolyl group substituted by 1 or 2 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms; an amino group and an acetamino group, an imidazo[2,1-b]thiazolyl

20 group, an imidazo[2,1-b]thiazolyl group substituted by 1 to 3 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms and a halogen atom, a phenylalkyl group having 7 to 10 carbon atoms, a quinolyl, a pyridyl, a naphthyl group, a naphthylalkyl


25 group having 11 to 15 carbon atoms, a dimethylamino-naphthyl group, a group represented by the formula:

5



wherein X is -O- or -S-,

a group represented by the formula:




- 5 a phenyl group, a phenyl group substituted by 1 to 5 members selected from the group consisting of a hydroxyl group; a methylsulfonyl group; an alkyl group having 1 to 6 carbon atoms; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group; a
- 10 dimethylamino group; an acetylamino group; a pyridyl group; a trifluoromethyl group; a trifluoromethoxy group and a halogen atom, a pyridyl group, a pyridyl group substituted by 1 or 2 members selected from the group consisting of a hydroxyl group; an alkyl group having 1
- 15 to 6 carbon atoms; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group; a dimethylamino group; an acetylamino group; a pyridyl group; a trifluoromethyl group; a trifluoromethoxy group and a halogen atom, a quinolyl group, or a quinolyl
- 20 group substituted by 1 or 2 members selected from the group consisting of a hydroxyl group; an alkyl group having 1 to 6 carbon atoms; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group; a dimethylamino group; an acetylamino group; a


pyridyl group; a trifluoromethyl group; a trifluoromethoxy group and a halogen atom, and  $R^{10}$  is a hydrogen atom or an alkyl group having 1 to 6 carbon atoms,

$R^2$  is a hydrogen atom, an alkyl group having 1 to 6 carbon atoms or a cinnamyl group,


$R^3$  is a group represented by the formula:




a group represented by the formula:



10 a group represented by the formula:



a group represented by the formula:



or a group represented by the formula:

15  $-\text{OCO}-\text{O}-\text{R}^{11}$

wherein  $R^{11}$  is a pyridylmethyl group, a methylthiomethyl group, a quinolyl group, a phenyl group, a phenyl group substituted by 1 to 5 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms;

20 a nitro group; an alkoxy group having 1 to 3 carbon atoms and a halogen atom, a pyridyl group, or a pyridyl group substituted by 1 or 2 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms; a nitro group; an alkoxy group having 1 to 3

25 carbon atoms and a halogen atom,

$R^4$  is a hydrogen atom, or  $R^3$  and  $R^4$  together form an oxo group, and

$R^5$  and  $R^6$  are the same or different, and are each a




hydrogen atom or an alkyl group having 1 to 6 carbon atoms, or a pharmaceutically acceptable salt thereof.

In the present invention, examples of the alkyl group having 1 to 6 carbon atoms are a methyl group, an ethyl group, a propyl group, a butyl group, a 3-methylbutyl group and a cyclohexyl group; examples of the alkoxy group having 1 to 3 carbon atoms are a methoxy group, an ethoxy group, a propoxy group and an isopropoxy group; and the halogen atom refers to a fluorine atom, a chlorine atom, a bromine atom and an iodine atom.

The pharmaceutically acceptable salt refers to a salt used in chemotherapy or prophylaxis of bacterially infectious diseases, for example, a salt with acetic acid, propionic acid, butyric acid, formic acid, trifluoroacetic acid, maleic acid, tartaric acid, citric acid, stearic acid, succinic acid, ethylsuccinic acid, lactobionic acid, gluconic acid, glucoheptonic acid, benzoic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, laurylsulfuric acid, malic acid, aspartic acid, glutaminic acid, adipic acid, cysteine, N-acetylcysteine, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, hydroiodic acid, nicotinic acid, oxalic acid, picric acid, thiocyanic acid, undecanoic acid, polyacrylate or carboxyvinyl polymer.

The compounds of the present invention can be

prepared, for example, according to the following reaction scheme.



Step (1); 2'-O-Acetyl-5-O-desosaminy-6-O-methylerythrionolide A described in US patent No.

5,523,399 is reacted with triphosgene in an inert solvent in the presence of pyridine under ice-cooling to  
5 give a compound of Formula (a). Examples of the inert solvent to be used here are dichloromethane, dichloroethane, acetone and tetrahydrofuran.

Step (2); The compound of Formula (a) is treated with a base in an inert solvent at a temperature  
10 of from room temperature to 120°C to give a compound of Formula (b). Examples of the inert solvent to be used here are N,N-dimethylformamide, dimethyl sulfoxide, N-methylpiperidone, tetrahydrofuran and a mixture thereof, and examples of the base to be used here are 1,1,3,3-  
15 tetramethylguanidine and potassium carbonate.

Step (3); The compound of Formula (b) is treated with a reagent of the following formula:




wherein  $R^{11}$  is as defined above, and an activating agent  
20 thereof in an inert solvent in the presence of a base such as 4-dimethylaminopyridine at a temperature of from -30°C to 30°C to give a compound of Formula (c).

Examples of the activating agent to be used herein are 1,3-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-  
25 3-ethylcarbodiimide hydrochloride and pivaloyl chloride. Examples of the inert solvent to be used here are dichloromethane, dichloroethane, acetone, pyridine, ethyl acetate and tetrahydrofuran.

Step (4); The compound of Formula (c) is reacted with 1,1'-carbonyldiimidazole in an inert solvent under the presence of a base such as sodium hydride to give a compound of Formula (d). The inert  
5 solvent is the same as used in Step (3).

Step (5); The compound of Formula (d) is reacted in an inert solvent with an amine compound of the following formula:



wherein  $n$ ,  $\text{R}^5$  and  $\text{R}^6$  are as defined above, and then is deprotected at the 2'-position by an ordinary methanolysis to give a compound of Formula (e). Examples of the inert solvent to be used here are  
20 acetonitrile, tetrahydrofuran, N,N-dimethylformamide, dioxane, ethyl acetate, N-methylpyrrolidone, a mixture of the solvent and water and a mixture thereof.

Step (6); The compound of Formula (e) is reacted with a reagent of the following formula:



wherein  $\text{R}^9$  is as defined above, in an inert solvent in the presence of a base such as pyridine, to give a compound of Formula (f) which is a compound of the present invention. The inert solvent is the same as  
30 used in Step (3).

The compounds of the present invention can be administered orally or parenterally in the dosage form

such as, for example, tablets, capsules, powders, troches, ointments, suspensions, suppositories and injections, all of which can be prepared according to conventional preparation techniques. The dose of the present compounds for treating an adult is from 100 to 1000 mg/day in single or several divided doses. This dose can be increased or decreased depending on the age, body weight and conditions of the patient.

#### 10 BEST MODE FOR CARRYING OUT THE INVENTION

The present invention is illustrated in more detail by the following Examples and a Test Example.

##### Reference Example 1

#### 15 Synthesis of 11-(2-aminoethyl)amino-11-deoxy-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

(1) To a solution of 70.0 g (77 mmol) of 10,11-anhydro-2',4"-di-O-acetyl-12-O-imidazolylcarbonyl-6-O-methylerythromycin A described in European Patent No. 638584 in 1 L of acetonitrile was added 30.0 ml (0.23 mol) of ethylenediamine at room temperature, followed by stirring overnight. The reaction solution was evaporated under reduced pressure, and the residue was dissolved in 1 L of methanol and refluxed under heating for 4 hours. After evaporation of the solvent, purification by silica gel column chromatography (chloroform : methanol : aqueous ammonia =20:1:0.1) gave

67.0 g (yield: 97 %) of 4"-O-acetyl-11-(2-aminoethyl)-amino-11-deoxy-6-O-methylerythromycin A 11,12-cyclic carbamate.

(2) A solution of 20 g (24 mmol) of the compound obtained in the above (1) in 100 ml of 1N aqueous hydrochloric acid solution was stirred at 70°C for an hour. The mixture was cooled to room temperature, and extracted with chloroform to remove the cladinose. The aqueous layer was made basic with an aqueous sodium hydroxide solution, extracted with chloroform and washed with a saturated aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The resulting residue was crystallized from 50 ml of ether to give 9.8 g (yield: 63 %) of the title compound. SIMS m/z: 658 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm): 2.25 (s, 6H, NMe<sub>2</sub>), 3.00 (s, 3H, 6-OMe), 4.40 (d, 1H, J=7.3 Hz, H-1'), 5.22 (dd, 1H, J=11.0, 2.4 Hz, H-13).

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm): 40.3 (NMe<sub>2</sub>), 49.6 (6-OMe), 106.9 (C1'), 157.9 (11,12-carbamate), 175.7 (C1), 215.6 (C9).

#### Example 1

11-[2-[(4-Acetaminophenyl)sulfonylamino]-ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate

(1) To a solution of 1.38 g (2.1 mmol) of the compound obtained in Reference Example 1 in 30 ml of a mixture of methylene chloride and pyridine was added 0.59 g (2.5 mmol) of p-acetamidobenzenesulfonyl chloride under ice-cooling, followed by stirring for 1.5 hours at room temperature. The reaction solution was diluted with ethyl acetate, and washed with an aqueous sodium hydroxide solution and a saturated aqueous sodium chloride solution respectively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give 2.07 g of 11-[2-[(4-acetaminophenyl)sulfonylamino]-ethyl]amino-11-deoxy-5-O-desosaminy-6-O-methyl-erythronolide A 11,12-cyclic carbamate.

(2) 2.07 g of the compound obtained in the above (1) was dissolved in 20 ml of acetone, and 0.3 ml (3.2 mmol) of acetic anhydride was added thereto at room temperature, followed by stirring overnight. After the reaction, the reaction solution was diluted with ethyl acetate, and washed with a saturated aqueous sodium bicarbonate solution and a saturated aqueous sodium chloride solution respectively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give 1.90 g of a 2'-O-acetyl compound.

(3) To a solution of 1.00 g (1.1 mmol) of the compound obtained in the above (2) in 20 ml of methylene chloride were successively added 0.58 g (3.3 mmol) of 2-

pyridylacetate hydrochloride, 0.64 g (3.3 mmol) of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and 0.14 g (1.1 mmol) of 4-dimethylaminopyridine under ice-cooling, followed by stirring at room temperature overnight. After the reaction, the reaction solution was diluted with ethyl acetate, and washed with an aqueous sodium hydroxide solution, a saturated aqueous ammonium chloride solution and a saturated aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give 1.06g of the residue, which was then dissolved in 20 ml of methanol and stirred at room temperature overnight. After the reaction, the solvent was evaporated under reduced pressure, and purification by silica gel column chromatography (chloroform : methanol : aqueous ammonia =15:1:0.1) gave 0.85 g (yield: 78 %) of the title compound.

IonSprayMS m/z: 974.5 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm): 2.03 (s, 3H, -COCH<sub>3</sub>), 2.28 (s, 6H, NMe<sub>2</sub>), 2.52 (s, 3H, 6-OMe), 3.95 and 3.98 (each d, each 1H, J<sub>gem</sub>=16.2 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.07 (d, 1H, J=7.3 Hz, H-1'), 4.91 (d, 1H, J=11.0 Hz, H-3), 4.92 (dd, 1H, J=11.2, 1.1 Hz, H-13), 5.49 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-), 8.14 (brs, 1H, -NHAc).

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm): 24.3 (-COCH<sub>3</sub>), 40.3 (NMe<sub>2</sub>), 49.7 (6-OMe), 103.4 (C1'), 157.5 (11,12-carbamate), 168.7 (-COCH<sub>3</sub>), 171.1 (-COCH<sub>2</sub>[2-



Pyridine)], 175.0 (C1), 215.7 (C9).

Example 2

11-[2-[(4-Acetaminophenyl)sulfonylamino]-  
5 ethylamino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-  
methylethrythrionolide A 11,12-cyclic carbamate

To a solution of 0.87 g (0.97 mmol) of the compound obtained in Example 1(2) in 18 ml of methylene chloride were successively added 0.69 ml (9.7 mmol) of dimethyl sulfoxide, 0.56 g (2.9 mmol) of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and 0.56 g (2.9 mmol) of pyridinium trifluoroacetate under ice-cooling, followed by stirring at room temperature overnight. After the reaction, the reaction solution  
15 was diluted with ethyl acetate, and washed with an aqueous sodium hydroxide solution and a saturated aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to  
20 give 0.99 g of the residue, which was then dissolved in 20 ml of methanol and stirred at room temperature overnight. After the reaction, the solvent was evaporated under reduced pressure, and purification by silica gel column chromatography (chloroform :  
25 methanol : aqueous ammonia =15:1:0.1) gave 0.55 g (yield: 66 %) of the title compound.

IonSprayMS m/z: 853.5 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.20 (s, 3H, -NHCOCH<sub>3</sub>),

2.26 (s, 6H, NMe<sub>2</sub>), 2.54 (s, 3H, 6-OMe), 3.83  
 (q, J=6.7 Hz, H-2), 4.22 (d, 1H, J=8.6 Hz, H-5),  
 4.27 (d, 1H, J=7.3 Hz, H-1'), 4.88 (dd, 1H, J=11.0,  
 2.4 Hz, H-13), 5.77 (brt, 1H, J=5.5 Hz, -NH<sub>2</sub>SO<sub>2</sub>-),  
 5 8.00 (s, 1H, -NHCOCH<sub>3</sub>)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 24.6 (-NHCOCH<sub>3</sub>), 40.2  
 (NMe<sub>2</sub>), 49.7 (6-OMe), 104.0 (C1'), 158.0 (11,12-  
 carbamate), 168.8 (-NHCOCH<sub>3</sub>), 170.1 (C1), 203.5 (C3),  
 216.4 (C9).

10

### Example 3

11-[2-[(2-Nitrophenyl)sulfonylamino]ethyl]-  
amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-  
O-methylerythronolide A 11,12-cyclic carbamate

15 (1) Following the same procedures as in  
 Example 1(1) and (2) using 1.57 g (2.4 mmol) of the  
 compound obtained in Reference Example 1 and 0.63 g (2.8  
 mmol) of 2-nitrobenzenesulfonyl chloride, there was  
 obtained 2.16 g of the 2'-O-acetyl compound.

20 (2) Following the same procedure as in Example  
 1(3) using 1.03 g (1.2 mmol) of the compound obtained in  
 the above (1), there was obtained 0.66 g (yield: 59 %) of  
 the title compound.

SIMS m/z: 962 [M+H]<sup>+</sup>

25 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.99  
 (s, 3H, 6-OMe), 3.92 and 3.96 (each d, each 1H,  
 J<sub>gem</sub>=16.2 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz,  
 H-1'), 5.02 (d, 1H, J=11.0 Hz, H-3), 5.06 (dd, 1H,

J=11.0, 2.1 Hz, H-13), 6.26 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.3  
(6-OMe), 103.5 (C1'), 157.5 (11,12-carbamate), 170.4  
(-COCH<sub>2</sub>[2-Pyridine]), 174.8 (C1), 215.8 (C9).

5

## Example 4

11-[2-[(2-Nitrophenyl)sulfonylamino]ethyl]-  
amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-  
methylethronolide A 11,12-cyclic carbamate

10

Following the same procedure as in Example 2  
using 1.06 g (1.2 mmol) of the compound obtained in  
Example 3(1), there was obtained 0.88 g (yield: 88 %) of  
the title compound.

IonSprayMS m/z: 841.4 [M+H]<sup>+</sup>

15

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ (ppm); 2.26 (s, 6H, NMe<sub>2</sub>), 2.58  
(s, 3H, 6-OMe), 3.84 (q, J=6.7 Hz, H-2), 4.24 (d, 1H,  
J=9.0 Hz, H-5), 4.27 (d, 1H, J=7.3 Hz, H-1'), 4.96  
(dd, 1H, J=10.8, 2.3 Hz, H-13)

20 Example 5

11-[2-[[1-(5-Dimethylamino)naphthyl]sulfonyl]-  
aminoethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-  
desosaminyl-6-O-methylethronolide A 11,12-cyclic  
carbamate

25

(1) Following the same procedures as in  
Example 1(1) and (2) using 1.44 g (2.2 mmol) of the  
compound obtained in Reference Example 1 and 0.71 g (2.6  
mmol) of 1-dimethylaminonaphthalene-5-sulfonyl chloride,

there was obtained 2.18 g of the 2'-O-acetyl compound.

(2) Following the same procedure as in Example 2(3) using 1.02 g (1.1 mmol) of the compound obtained in the above (1), there was obtained 0.95 g (yield: 86 %) of the title compound.

IonSprayMS m/z: 1010.6 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (300MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.71 (s, 3H, 6-OMe), 2.86 (s, 6H, [Naphthalene]-NMe<sub>2</sub>), 3.95 (s, 2H, -CH<sub>2</sub>[2-Pyridine]), 4.05 (d, 1H, J=7.3 Hz, H-1'), 4.99 - 5.06 (m, 2H, H-3 and H-13), 6.23 (brt, 1H, J=6.0 Hz, -NH<sub>2</sub>SO<sub>2</sub>-).

#### Example 6

11-[2-[[1-(5-Dimethylamino)naphthyl]sulfonyl-  
amino]ethyl]amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-  
O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 2 using 1.19 g (1.3 mmol) of the compound obtained in Example 5(1), there was obtained 0.91 g (yield: 79 %) of the title compound.

SIMS m/z: 889 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.26 (s, 6H, NMe<sub>2</sub>), 2.44 (s, 3H, 6-OMe), 2.88 (s, 6H, [Naphthalene]-NMe<sub>2</sub>), 3.83 (q, J=6.7Hz, H-2), 4.20 (d, 1H, J=8.5Hz, H-5), 4.27 (d, 1H, J=7.3 Hz, H-1'), 4.91 (dd, 1H, J=11.0, 2.4 Hz, H-13), 6.11 (brt, 1H, J=5.5Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.2 (NMe<sub>2</sub>), 45.4 ([Naphthalene]-NMe<sub>2</sub>), 49.6 (6-OMe), 103.9 (C1'),

157.9 (11,12-carbamate), 170.0 (C1), 203.4 (C3),  
216.3 (C9).

Example 7

5            11-[2-(Benzo-2,1,3-thiadiazole-4-sulfonyl-  
amino)ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-  
desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
carbamate

(1) Following the same procedures as in  
10 Example 1(1) and (2) using 1.21 g (1.8 mmol) of the  
compound obtained in Reference Example 1 and 0.52 g (2.2  
mmol) of benzo-2,1,3-thiadiazole-4-sulfonyl chloride,  
there was obtained 1.75 g of the 2'-O-acetyl compound.

(2) Following the same procedure as in Example  
15 1(3) using 0.85 g (0.95 mmol) of the compound obtained  
in the above (1), there was obtained 0.78 g (yield:  
85 %) of the title compound.

SIMS m/z: 975 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.91  
20        (s, 3H, 6-OMe), 3.92 and 3.98 (each d, each 1H,  
J<sub>gem</sub>=16.1 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.4 Hz,  
H-1'), 5.01 (d, 1H, J=11.2 Hz, H-3), 5.04 (dd, 1H,  
J=11.0, 2.1 Hz, H-13), 6.28 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.2  
25        (6-OMe), 103.6 (C1'), 157.3 (11,12-carbamate), 170.4  
(-COCH<sub>2</sub>[2-Pyridine]), 174.8 (C1), 215.7 (C9).

## Example 8

11-[2-(Benzo-2,1,3-thiadiazole-4-sulfonylamino)ethylamino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
5 carbamate

Following the same procedure as in Example 2 using 0.85 g (0.94 mmol) of the compound obtained in Example 7(1), there was obtained 0.67 g (yield: 84 %) of the title compound.

10 SIMS m/z: 854 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.26 (s, 6H, NMe<sub>2</sub>), 2.51 (s, 3H, 6-OMe), 3.83 (q, J=6.7 Hz, H-2), 4.22 (d, 1H, J=9.2 Hz, H-5), 4.27 (d, 1H, J=7.3 Hz, H-1'), 4.93 (dd, 1H, J=11.0, 2.4 Hz, H-13), 6.15 (brs, 1H,  
15 -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 24.6 (-NHCOCH<sub>3</sub>), 40.2 (NMe<sub>2</sub>), 49.8 (6-OMe), 104.0 (C1'), 157.3 (11,12-carbamate), 170.0 (C1), 203.4 (C3), 216.3 (C9).

## 20 Example 9

11-[2-[(8-Quinolyl)sulfonylamino]ethylamino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythro-  
nolide A 11,12-cyclic carbamate

(1) Following the same procedures as in  
25 Example 1(1) and (2) using 2.02 g (3.1 mmol) of the compound obtained in Reference Example 1 and 0.84 g (3.7 mmol) of quinoline-8-sulfonyl chloride, there was obtained 3.08 g of the 2'-O-acetyl compound.

(2) Following the same procedure as in Example 2 using 3.08 g of the compound obtained in the above (1), there was obtained 2.45 g (yield: 94 %) of the title compound.

5 IonSpray MS m/z: 847.4 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.26 (s, 6H, NMe<sub>2</sub>), 2.43 (s, 3H, 6-OMe), 3.80 (q, J=7.0 Hz, H-2), 4.18 (d, 1H, J=8.8 Hz, H-5), 4.27 (d, 1H, J=7.4 Hz, H-1'), 4.93 (dd, 1H, J=10.6, 2.4 Hz, H-13), 6.73 (brt, 1H, J=6.4 Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

10 <sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.2 (NMe<sub>2</sub>), 49.6 (6-OMe), 104.0 (C1'), 157.4 (11,12-carbamate), 169.4 (C1), 203.8 (C3), 216.0 (C9).

15 Example 10

11-[2-[(8-Quinolyl)sulfonylamino]ethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

(1) To a solution of 53.6 g (0.09 mol) of 2'-  
20 O-acetyl-5-O-desosaminyl-6-O-methylerythronolide A in 500 ml of methylene chloride was added 54.9 ml (0.70 mol) of pyridine. 16.8 g (0.06 mol) of triphosgene was added to the mixture under ice-cooling, followed by stirring at room temperature for 2 hours. After the  
25 reaction, water was added to the reaction mixture under ice-cooling to decompose the excess triphosgene, and the mixture was diluted with chloroform and washed with water and a saturated aqueous sodium chloride solution

successively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give 56.0 g of the 11,12-cyclic carbonate compound.

5                   (2) To a solution of 56.0 g of the compound obtained in the above (1) in 500 ml of N,N-dimethylformamide was added 23.7 ml (0.23 mol) of 1,1,3,3-tetramethylguanidine, followed by stirring at 100°C for 3 hours. After the reaction, the reaction  
10 solution was diluted with ethyl acetate and washed with distilled water and a saturated aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give 50.27 g of  
15 10,11-anhydro-2'-O-acetyl-5-O-desosaminy-6-O-methylerythronolide A.

(3) Following the same procedure as in Example 2(3) using 50.3 g (0.08 mol) of the compound obtained in the above (2), there was obtained 41.95 g of 10,11-anhydro-2'-O-acetyl-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A.  
20

(4) To a solution of 31.0 g (0.04 mol) of the compound obtained in the above (3) in a mixture of 180 ml of N,N-dimethylformamide and 120 ml of  
25 tetrahydrofuran were successively added 20.6 g (0.13 mol) of carbonyldiimidazole and 3.38 g (0.08 mol) of sodium hydride under ice-cooling, followed by stirring under ice-cooling for 40 minutes. After the reaction,



the reaction solution was diluted with ethyl acetate and washed with distilled water and a saturated aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, and  
5 the solvent was evaporated under reduced pressure to give 32.7 g of 10,11-anhydro-2'-O-acetyl-12-O-imidazolylcarbonyl-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A.

(5) To a solution of 11.5 g (14.6 mmol) of the  
10 compound obtained in the above (4) in 100 ml of acetonitrile was added 9.8 ml (0.15 mol) of ethylenediamine, followed by stirring for 2 days. After the reaction, the reaction solution was diluted with ethyl acetate and washed with distilled water and a  
15 saturated aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The resulting residue was dissolved in 100 ml of methanol, followed by stirring at room temperature  
20 overnight. After the reaction, the reaction solution was evaporated under reduced pressure, and the precipitated crystals were washed with ether to give 3.98 g of 11-(2-aminoethyl)amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A  
25 11,12-cyclic carbamate.

(6) To a solution of 0.62 g (0.8 mmol) of the compound obtained in the above (5) in 10 ml of a mixture of methylene chloride and pyridine was added 0.22 g (1.0

mmol) of quinoline-8-sulfonyl chloride under ice-cooling, followed by stirring at room temperature for 2 hours. After the reaction, the reaction solution was diluted with ethyl acetate and washed with an aqueous sodium hydroxide solution and a saturated aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure, and purification by silica gel column chromatography (chloroform : methanol : aqueous ammonia =20:1:0.1) gave 0.62 g (yield: 80 %) of the title compound.

IonSprayMS m/z: 968.5 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.81 (s, 3H, 6-OMe), 3.93 and 3.97 (each d, each 1H, J<sub>gem</sub>=16.2 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'), 5.04 (d, 1H, J=11.0 Hz, H-3), 5.11 (dd, 1H, J=11.0, 2.4 Hz, H-13), 6.79 (brt, 1H, J=6.4 Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 49.9 (6-OMe), 103.6 (C1'), 157.5 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 174.1 (C1), 215.7 (C9).

#### Example 11

11-[2-[(3-Nitrophenyl)sulfonylamino]ethyl]-amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.53 g (0.68 mmol) of the compound obtained

in Example 10(5) and 0.18 g (0.23 mmol) of 3-nitro-benzenesulfonyl chloride, there was obtained 0.55 g (yield: 85 %) of the title compound.

SIMS m/z: 962 [M+H]<sup>+</sup>

- 5 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.87 (s, 3H, 6-OMe), 3.93 and 3.96 (each d, each 1H, J<sub>gem</sub>=16.2 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'), 4.92 (dd, 1H, J=11.0, 2.4Hz, H-13), 4.99 (d, 1H, J=11.0 Hz, H-3), 6.31 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)
- 10 <sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.1 (6-OMe), 103.5 (C1'), 157.7 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 175.5 (C1), 216.0 (C9).

#### Example 12

- 15 11-[2-[(3-Nitrophenyl)sulfonylamino]ethyl]-amino-3,11-dideoxy-3-oxo-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate

(1) Following the same procedure as in Example 1 for oxidation using 72.0 g of the compound obtained in  
20 Example 10(2), there was obtained 67.0 g of 2'-O-acetyl-3,11-dideoxy-3-oxo-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbonate.

(2) Following the same procedure as in Example 10(2) using 67.0 g of the compound obtained in the above  
25 (1), there was obtained 19.8 g of the 10,11-anhydro compound.

(3) Following the same procedure as in Example 10(4) using 19.8 g of the compound obtained in the above

(2), purification by silica gel column chromatography (acetone : n-hexane : triethylamine =10:20:0.2) gave 15.5 g (yield: 68 %) of the 12-O-imidazolylcarbonyl compound.

5                   (4) To a solution of 8.4 g (12 mmol) of the compound obtained in the above (3) in 60 ml of acetonitrile was added 8.0 ml (0.12 mol) of ethylenediamine, followed by stirring at room temperature for 2 days. After the reaction, the reaction solution was diluted  
10 with ethyl acetate and washed with distilled water and a saturated aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure to give 7.1 g of 2'-O-acetyl-11-(2-aminoethyl)-  
15 amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methyle-  
erythronolide A 11,12-cyclic carbamate.

(5) To a solution of 0.52 g (0.75 mmol) of the compound obtained in the above (4) in 10 ml of a mixture of methylene chloride and pyridine was added 0.20 g (0.9  
20 mmol) of 3-nitrobenzenesulfonyl chloride under ice-cooling, followed by stirring at room temperature overnight. After the reaction, the reaction solution was diluted with ethyl acetate and washed with an aqueous sodium hydroxide solution and a saturated  
25 aqueous sodium chloride solution successively. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was dissolved in 20 ml of methanol, followed

by stirring at room temperature overnight. After the reaction, the solvent was evaporated under reduced pressure, and purification by silica gel column chromatography (chloroform : methanol : aqueous ammonia  
5 =27:1:0.1) gave 0.26 g (yield: 41 %) of the title compound.

SIMS m/z: 841 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.26 (s, 6H, NMe<sub>2</sub>), 2.49  
(s, 3H, 6-OMe), 3.83 (q, J=7.0 Hz, H-2), 4.21 (d, 1H,  
10 J=8.8 Hz, H-5), 4.26 (d, 1H, J=7.3 Hz, H-1'), 4.82  
(dd, 1H, J=10.9, 2.4 Hz, H-13), 6.12 (brs, 1H,  
-NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.2 (NMe<sub>2</sub>), 50.0  
(6-OMe), 104.0 (C1'), 158.0 (11,12-carbamate), 170.4  
15 (C1), 203.1 (C3), 216.6 (C9).

#### Example 13

11-[2-[(4-Nitrophenyl)sulfonylamino]ethyl]-  
amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-  
20 O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example  
10(6) using 0.51 g (0.66 mmol) of the compound obtained  
in Example 10(5) and 0.17 g (0.77 mmol) of 4-nitro-  
benzenesulfonyl chloride, there was obtained 0.51 g  
25 (yield: 80 %) of the title compound.

SIMS m/z: 962 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.89  
(s, 3H, 6-OMe), 3.93 and 3.96 (each d, each 1H,

$J_{gem}=16.5\text{Hz}$ ,  $-\text{CH}_2[2\text{-Pyridine}]$ ), 4.06(d, 1H,  $J=7.3\text{ Hz}$ , H-1'), 4.91 (dd, 1H,  $J=11.0, 1.8\text{ Hz}$ , H-13), 5.00 (d, 1H,  $J=11.0\text{ Hz}$ , H-3), 6.32 (brs, 1H,  $-\text{NH}\text{SO}_2-$ )  
 $^{13}\text{C}$ -NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 40.2 ( $\text{NMe}_2$ ), 50.1  
 5 (6-OMe), 103.5 ( $\text{C1}'$ ), 157.9 (11,12-carbamate), 170.4 ( $-\text{COCH}_2[2\text{-Pyridine}]$ ), 175.3 (C1), 216.1 (C9).

## Example 14

11-[2-[(4-Nitrophenyl)sulfonylamino]ethyl]-  
 10 amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-  
methylethronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 12(5) using 0.83 g (1.2 mmol) of the compound obtained in Example 12(4) and 0.35 g (1.6 mmol) of 4-nitro-  
 15 benzenesulfonyl chloride, there was obtained 0.41 g (yield: 41 %) of the title compound.

SIMS  $m/z$ : 841  $[\text{M}+\text{H}]^+$

$^1\text{H}$ -NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 2.26 (s, 6H,  $\text{NMe}_2$ ), 2.51 (s, 3H, 6-OMe), 3.83 (q,  $J=6.7\text{ Hz}$ , H-2), 4.22 (d, 1H,  $J=9.2\text{ Hz}$ , H-5), 4.27 (d, 1H,  $J=7.3\text{ Hz}$ , H-1'), 4.82 (dd, 1H,  $J=11.0, 2.4\text{ Hz}$ , H-13), 6.14 (brs, 1H,  $-\text{NH}\text{SO}_2-$ )

$^{13}\text{C}$ -NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 40.2 ( $\text{NMe}_2$ ), 49.7 (6-OMe), 104.0 ( $\text{C1}'$ ), 158.2 (11,12-carbamate), 170.3 (C1), 203.2 (C3), 216.7 (C9).

## Example 15

11-[2-(Pentafluorophenyl)sulfonylamino]-

ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-  
desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
carbamate

Following the same procedure as in Example

- 5 10(6) using 0.53 g (0.68 mmol) of the compound obtained in Example 10(5) and 0.22 g (0.83 mmol) of pentafluorobenzenesulfonyl chloride, there was obtained 0.29 g (yield: 42 %) of the title compound.

SIMS m/z: 1007 [M+H]<sup>+</sup>

- 10 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>),  
3.04 (s, 3H, 6-OMe), 3.78 (d, 1H, J=4.3 Hz, H-5),  
3.92 and 3.96 (each d, each 1H, J<sub>gem</sub>=15.9 Hz,  
-CH<sub>2</sub>[2-Pyridine]), 4.07 (d, 1H, J=7.3 Hz, H-1'),  
4.99 (dd, 1H, J=11.0, 2.4 Hz, H-13), 5.03 (d, 1H,  
15 J=11.0 Hz, H-3)  
<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.4  
(6-OMe), 103.5 (C1'), 157.7 (11,12-carbamate), 170.5  
(-COCH<sub>2</sub>[2-Pyridine]), 175.5 (C1), 216.0 (C9).

20 Example 16

11-[2-(Pentafluorophenylsulfonylamino)-  
ethylamino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-  
methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example

- 25 12(5) using 0.83 g (1.2 mmol) of the compound obtained in Example 12(4) and 0.38 g (1.4 mmol) of pentafluorobenzenesulfonyl chloride, there was obtained 0.49 g (yield: 46 %) of the title compound.

SIMS m/z: 886 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.26 (s, 6H, NMe<sub>2</sub>), 2.62 (s, 3H, 6-OMe), 3.86 (q, J=6.7 Hz, H-2), 4.25 (d, 1H, J=9.2 Hz, H-5), 4.27 (d, 1H, J=7.3 Hz, H-1'), 4.90 (dd, 1H, J=11.0, 2.4 Hz, H-13)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.2 (NMe<sub>2</sub>), 49.9 (6-OMe), 104.0 (C1'), 157.7 (11,12-carbamate), 170.7 (C1), 203.0 (C3), 216.5 (C9).

10 Example 17

11-[2-(Methanesulfonylamino)ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example

15 10(6) using 0.51 g (0.66 mmol) of the compound obtained in Example 10(5) and 0.06 ml (0.79 mmol) of methanesulfonyl chloride, there was obtained 0.45 g (yield: 80 %) of the title compound.

SIMS m/z: 855 [M+H]<sup>+</sup>

20 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.98 (s, 3H, -SO<sub>2</sub>Me), 3.05 (s, 3H, 6-OMe), 3.92 and 3.96 (each d, each 1H, J<sub>gem</sub>=15.9 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'), 5.03 (d, 2H, J=10.9 Hz, H-3 and H-13), 5.51 (brt, 1H, J=5.5 Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

25 <sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 41.2 (-SO<sub>2</sub>Me), 50.3 (6-OMe), 103.5 (C1'), 157.9 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 175.1 (C1), 216.0 (C9).



## Example 18

11-[2-(methanesulfonylamino)ethylamino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

5           Following the same procedure as in Example 12(5) using 0.83 g (1.2 mmol) of the compound obtained in Example 12(4) and 0.11 ml (1.4 mmol) of methanesulfonyl chloride, there was obtained 0.18 g (yield: 21 %) of the title compound.

10   FABMS m/z: 734 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.27 (s, 6H, NMe<sub>2</sub>), 2.66 (s, 3H, 6-OMe), 2.99 (s, 3H, -SO<sub>2</sub>Me), 3.86 (q, J=6.7 Hz, H-2), 4.25 (d, 1H, J=9.2 Hz, H-5), 4.28 (d, 1H, J=7.3 Hz, H-1'), 4.95 (dd, 1H, J=11.0, 2.4 Hz, H-13),  
15   5.33 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.2 (NMe<sub>2</sub>), 41.0 (-SO<sub>2</sub>Me), 49.9 (6-OMe), 104.0 (C1'), 157.9 (11,12-carbamate), 170.3 (C1), 203.3 (C3), 216.5 (C9).

## 20   Example 19

11-[2-(Phenylsulfonylamino)ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

          Following the same procedure as in Example  
25   10(6) using 0.54 g (0.69 mmol) of the compound obtained in Example 10(5) and 0.15 g (0.85 mmol) of benzene-sulfonyl chloride, there was obtained 0.52 g (yield: 82 %) of the title compound.

IonSpray MS m/z: 917.4 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.28 (s, 6H, NMe<sub>2</sub>), 2.80  
(s, 3H, 6-OMe), 3.92 and 3.96 (each d, each 1H,  
J<sub>gem</sub>=15.8 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.05 (d, 1H, J=7.3 Hz,  
5 H-1'), 4.95 (dd, 1H, J=11.0, 2.2 Hz, H-13), 4.99  
(d, 1H, J=11.0 Hz, H-3), 5.93 (brt, 1H, J=5.8 Hz,  
-NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.1  
(6-OMe), 103.5 (C1'), 157.7 (11,12-carbamate), 170.4  
10 (-COCH<sub>2</sub>[2-Pyridine]), 175.1 (C1), 215.9 (C9).

#### Example 20

11-[2-(1-Naphthalenesulfonylamino)ethylamino]-  
11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-  
15 methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example  
10(6) using 0.54 g (0.69 mmol) of the compound obtained  
in Example 10(5) and 0.19 g (0.84 mmol) of  
naphthalenesulfonyl chloride, there was obtained 0.58 g  
20 (yield: 87 %) of the title compound.

IonSpray MS m/z: 967.4 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.28 (s, 6H, NMe<sub>2</sub>), 2.67  
(s, 3H, 6-OMe), 3.93 and 3.97 (each d, each 1H,  
J<sub>gem</sub>=16.2 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.04 (d, 1H, J=7.3 Hz,  
25 H-1'), 5.02 (dd, 1H, J=11.2, 2.2 Hz, H-13), 5.00  
(d, 1H, J=11.0 Hz, H-3), 6.31 (brt, 1H, J=5.8 Hz,  
-NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.0

(6-OMe), 103.6 (C1'), 157.7 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 175.1 (C1), 215.8 (C9).

Example 21

5                    11-[2-(2-Mesitylenesulfonylamino)ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.53 g (0.68 mmol) of the compound obtained in Example 10(5) and 0.18 g (0.82 mmol) of 2-mesitylenesulfonyl chloride, there was obtained 0.55 g (yield: 84 %) of the title compound.

IonSpray MS m/z: 959.5 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.27 and 2.63 (each s, 9H, -PhMe<sub>3</sub>), 2.29 (s, 6H, NMe<sub>2</sub>), 2.93 (s, 3H, 6-OMe), 3.92 and 3.95 (each d, each 1H, J<sub>gem</sub>=16.2 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.05 (d, 1H, J=7.3 Hz, H-1'), 5.00-5.03 (m, 2H, H-3 and H-13), 5.92 (brt, 1H, J=6.0 Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

20    <sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 20.8 and 22.9 (-PhMe<sub>3</sub>), 40.3 (NMe<sub>2</sub>), 50.2 (6-OMe), 103.5 (C1'), 157.8 (11, 12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 174.8 (C1), 215.8 (C9).

25    Example 22

11-[2-[2-(1-Naphthyl)ethanesulfonylamino]-ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic

carbamate

Following the same procedure as in Example 10(6) using 0.53 g (0.68 mmol) of the compound obtained in Example 10(5) and 0.21 g (0.82 mmol) of 2-(1-naphthyl)ethanesulfonyl chloride, there was obtained 0.38 g (yield: 56 %) of the title compound.

IonSpray MS m/z: 995.5 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.31 (s, 6H, NMe<sub>2</sub>), 3.04 (s, 3H, 6-OMe), 3.91 and 3.96 (each d, each 1H, J<sub>gem</sub>=15.9 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'), 5.03 (d, 1H, J=11.0 Hz, H-3), 5.04 (dd, 1H, J=11.0, 2.2 Hz, H-13), 5.60 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.3 (6-OMe), 103.5 (C1'), 158.0 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 175.0 (C1), 216.0 (C9).

## Example 23

11-[2-(2-Acetamido-4-methyl-5-thiazolesulfonyl-amino)ethyl]lamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.57 g (0.73 mmol) of the compound obtained in Example 10(5) and 0.22 g (0.86 mmol) of 2-acetamido-4-methyl-5-thiazolesulfonyl chloride, recrystallization from ethyl acetate/n-hexane gave 0.19 g (yield: 26 %) of the title compound as the first crystals.

FABMS m/z: 995 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500MHz, CDCl<sub>3</sub>) δ (ppm); 2.21 (s, 3H, -NHCOCH<sub>3</sub>),  
 2.29 (s, 6H, NMe<sub>2</sub>), 2.49 (s, 3H, [Thiazole]-Me), 2.91  
 (s, 3H, 6-OMe), 3.91 and 3.95 (each d, each 1H,  
 J<sub>gem</sub>=15.8 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.05 (d, 1H, J=7.4 Hz,  
 5 H-1'), 4.97 (d, 2H, J=11.3 Hz, H-3 and H-13), 6.08  
 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 16.3 ([Thiazole]-Me),  
 23.0 (-NHCOCH<sub>3</sub>), 40.3 (NMe<sub>2</sub>), 50.2 (6-OMe), 103.6  
 (C1'), 157.8 (11,12-carbamate), 168.1 (-NHCOCH<sub>3</sub>),  
 10 170.5 (-COCH<sub>2</sub>[2-Pyridine]), 175.1 (C1), 215.9 (C9).

## Example 24

11-[2-[3,5-Dimethylisoxazole-4-sulfonyl-  
 amino]ethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-  
 15 desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
 carbamate

Following the same procedure as in Example  
 10(6) using 0.54 g (0.69 mmol) of the compound obtained  
 in Example 10(5) and 0.16 g (0.82 mmol) of 3,5-  
 20 dimethylisoxazole-4-sulfonyl chloride, there was  
 obtained 0.28 g (yield: 44 %) of the title compound.

FABMS m/z: 936 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.41  
 and 2.64 (each s, each 3H, [Isoxazole]-Me), 2.96 (s,  
 25 3H, 6-OMe), 3.92 and 3.96 (each d, each 1H, J<sub>gem</sub>=16.2  
 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'),  
 4.95 (dd, 1H, J=11.0, 2.1 Hz, H-13), 5.02 (d, 1H,  
 J=11.0 Hz, H-3), 6.31 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)

$^{13}\text{C}$ -NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 10.7 and 12.6  
 ([Isoxazole]-Me), 40.3 ( $\text{NMe}_2$ ), 50.1 (6-OMe),  
 103.6 ( $\text{C1}'$ ), 158.1 (11,12-carbamate), 170.5  
 ( $-\text{COCH}_2[2\text{-Pyridine}]$ ), 175.2 ( $\text{C1}$ ), 216.1 ( $\text{C9}$ ).

5

## Example 25

11-[2-[2-(Pyrid-2-yl)thiophene-5-  
sulfonylamino]ethylamino-11-deoxy-3-O-(2-pyridyl)-  
acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-  
 10 cyclic carbamate

Following the same procedure as in Example  
 10(6) using 0.53 g (0.69 mmol) of the compound obtained  
 in Example 10(5) and 0.21 g (0.81 mmol) of 2-(pyrid-2-  
 yl)thiophene-5-sulfonyl chloride, there was obtained  
 15 0.47 g (yield: 68 %) of the title compound.

FABMS  $m/z$ : 1000  $[\text{M}+\text{H}]^+$

$^1\text{H}$ -NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 2.29 (s, 6H,  $\text{NMe}_2$ ), 2.94  
 (s, 3H, 6-OMe), 3.92 and 3.96 (each d, each 1H,  
 $J_{\text{gem}}=15.9$  Hz,  $-\text{CH}_2[2\text{-Pyridine}]$ ), 4.05 (d, 1H,  $J=7.3$  Hz,  
 20 H-1'), 5.01 (dd, 1H,  $J=11.0$ , 2.4 Hz, H-13), 5.02  
 (d, 1H,  $J=11.0$  Hz, H-3), 6.14 (brs, 1H,  $-\text{NH}_2\text{SO}_2-$ )

$^{13}\text{C}$ -NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 40.3 ( $\text{NMe}_2$ ), 50.2  
 (6-OMe), 103.5 ( $\text{C1}'$ ), 157.8 (11,12-carbamate), 170.4  
 ( $-\text{COCH}_2[2\text{-Pyridine}]$ ), 175.0 ( $\text{C1}$ ), 215.9 ( $\text{C9}$ ).

25

## Example 26

11-[2-[(3-Pyridyl)sulfonylamino]ethylamino-11-  
deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methyl-

erythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 1.0 g (1.3 mmol) of the compound obtained in Example 10(5) and 0.69 g of pyridine-3-sulfonyl chloride described in the following Reference Example 2, there was obtained 0.20 g (yield: 17 %) of the title compound. IonSprayMS m/z: 918.4 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.86 (s, 3H, 6-OMe), 3.93 and 3.96 (each d, each 1H, J<sub>gem</sub>=15.9 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'), 4.93 (dd, 1H, J=11.0, 2.4 Hz, H-13), 5.00 (d, 1H, J=11.0 Hz, H-3), 6.21 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.3 (NMe<sub>2</sub>), 50.1 (6-OMe), 103.5 (C1'), 157.8 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 175.3 (C1), 216.0 (C9).

## Reference Example 2

Preparation of pyridine-3-sulfonyl chloride

1.6 g (0.01 mol) of pyridine-3-sulfonic acid and 4.2 g (0.02 mol) of phosphorus pentachloride together were stirred at 200°C for 2.5 hours. After the reaction, chloroform was added to the reaction system, the insoluble matter was removed by filtration, and the filtrate was evaporated under reduced pressure to give 2.41 g of pyridine-3-sulfonyl chloride.

## Example 27

11-[2-[(4-Methylphenyl)sulfonylamino]ethyl]-

amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.53 g (0.69 mmol) of the compound obtained in Example 10(5) and 0.15 g (0.80 mmol) of p-toluene-sulfonyl chloride, there was obtained 0.40 g (yield: 63 %) of the title compound.

IonSprayMS m/z: 931.5 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.39 (s, 3H, Ph-Me), 2.83 (s, 3H, 6-OMe), 3.93 and 3.96 (each d, each 1H, J<sub>gem</sub>=15.9 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'), 4.96 (dd, 1H, J=11.0, 2.4 Hz, H-13), 5.00 (d, 1H, J=11.0 Hz, H-3), 5.83 (brt, 1H, J=6.1 Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 21.4 (Ph-Me), 40.3 (NMe<sub>2</sub>), 50.0 (6-OMe), 103.5 (C1'), 157.7 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 175.0 (C1), 215.9 (C9).

Example 28

11-[2-[4-(4-Dimethylaminophenylazo)phenyl-sulfonylamino]ethylamino-11-deoxy-3-O-(2-pyridyl)-acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.52 g (0.67 mmol) of the compound obtained in Example 10(5) and 0.26 g (0.80 mmol) of 4-(4-dimethylaminophenylazo)benzenesulfonyl chloride, there



was obtained 0.54 g (yield: 76 %) of the title compound.

IonSprayMS m/z: 1064.6 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.93  
 (s, 3H, 6-OMe), 3.11 (s, 6H, Ph-NMe<sub>2</sub>), 3.92 and 3.96  
 5 (each d, each 1H, J=15.9 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.05  
 (d, 1H, J=7.3 Hz, H-1'), 4.98 (dd, J=11.0, 1.8 Hz,  
 H-13), 5.04 (d, 1H, J=11.0 Hz, H-3), 6.03 (brt, 1H,  
 J=6.1 Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>) δ (ppm); 40.2 (NMe<sub>2</sub>), 40.3 (NMe<sub>2</sub>),  
 10 50.2 (6-OMe), 103.5 (C1'), 157.9 (11,12-carbamate),  
 170.3 (-COCH<sub>2</sub>[2-Pyridine]), 174.9 (C1), 215.9 (C9).

#### Example 29

11-[2-[(4-Methoxyphenyl)sulfonylamino]ethyl]-  
 15 amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-  
O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example  
 10(6) using 0.57 g (0.74 mmol) of the compound obtained  
 in Example 10(5) and 0.18 g (0.87 mmol) of 4-methoxy-  
 20 benzenesulfonyl chloride, there was obtained 0.45 g  
 (yield: 64 %) of the title compound.

FABMS m/z: 947 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.83  
 (s, 3H, 6-OMe), 3.83 (s, 3H, Ph-OMe), 3.92 and 3.96  
 25 (each d, each 1H, J=15.9 Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06  
 (d, 1H, J=7.3 Hz, H-1'), 4.96 (dd, 1H, J=11.0, 1.8  
 Hz, H-13), 5.01 (d, 1H, J=11.0 Hz, H-3), 5.79 (brt,  
 1H, J=6.1 Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

$^{13}\text{C}$ -NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 40.3 ( $\text{NMe}_2$ ), 50.1 (6-OMe), 55.5 (Ph-OMe), 103.5 ( $\text{C1}'$ ), 157.7 (11,12-carbamate), 170.4 ( $-\text{COCH}_2[2\text{-Pyridine}]$ ), 175.0 ( $\text{C1}$ ), 215.8 ( $\text{C9}$ ).

5

## Example 30

11-[2-[(4-Cyanophenyl)sulfonylamino]ethyl]-amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

10

Following the same procedure as in Example 10(6) using 0.54 g (0.69 mmol) of the compound obtained in Example 10(5) and 0.17 g (0.84 mmol) of 4-cyanobenzenesulfonyl chloride, there was obtained 0.48 g (yield: 74 %) of the title compound.

15 FABMS  $m/z$ : 942  $[\text{M}+\text{H}]^+$ 

$^1\text{H}$ -NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm); 2.29 (s, 6H,  $\text{NMe}_2$ ), 2.86 (s, 3H, 6-OMe), 3.95 (s, 2H,  $-\text{CH}_2[2\text{-Pyridine}]$ ), 4.07(d, 1H,  $J=7.3$  Hz, H-1'), 4.91 (dd, 1H,  $J=11.0$ , 2.1 Hz, H-13), 5.01 (d, 1H,  $J=11.2$  Hz, H-3).

20

## Example 31

11-[2-[(4-Trifluoromethoxyphenyl)sulfonyl-amino]ethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

25

Following the same procedure as in Example 10(6) using 0.65 g (0.84 mmol) of the compound obtained in Example 10(5) and 0.26 g (1.0 mmol) of 4-(trifluoro-

methoxy)benzenesulfonyl chloride, there was obtained 0.36 g (yield: 43 %) of the title compound.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.84 (s, 3H, 6-OMe), 3.94 (s, 2H, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.1 Hz, H-1'), 4.93 (dd, 1H, J=11.0, 1.9 Hz, H-13), 5.01 (d, 1H, J=10.8 Hz, H-3).

#### Example 32

11-[2-[(4-Trifluoromethylphenyl)sulfonyl-aminolethyllamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.52 g (0.67 mmol) of the compound obtained in Example 10(5) and 0.20 g (0.82 mmol) of 4-(trifluoromethyl)benzenesulfonyl chloride, there was obtained 0.45 g (yield: 68 %) of the title compound.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ (ppm); 2.29 (s, 6H, NMe<sub>2</sub>), 2.85 (s, 3H, 6-OMe), 3.94 (s, 2H, -CH<sub>2</sub>[2-Pyridine]), 4.06 (d, 1H, J=7.3 Hz, H-1'), 4.91 (dd, 1H, J=11.0, 2.1 Hz, H-13), 5.01 (d, 1H, J=11.2 Hz, H-3).

#### Example 33

11-[2-(N-Phenylsulfonyl)ethyllamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

To a solution of 0.15 g (0.17 mmol) of the compound obtained in Example 10(4) in 10 ml of a mixture

of N,N-dimethylformamide and distilled water [10:3] was added 0.34 g (1.7 mmol) of 2-aminoethylsulfonanilide described in the following Reference Example 3 at room temperature, followed by stirring for 7 days. After the  
5 reaction, chloroform was added to the reaction solution, and the mixture was successively washed with distilled water and a saturated aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced  
10 pressure. The residue was dissolved in 10 ml of methanol and stirred at room temperature overnight. After the reaction, the solvent was evaporated under reduced pressure, and purification by silica gel column chromatography (chloroform : methanol : aqueous ammonia  
15 =19:1:0.1 - 9:1:0.1) gave 0.05 g (yield: 32 %) of the title compound.

FABMS m/z: 917 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500MHz, CDCl<sub>3</sub>) δ (ppm); 0.84 (t, 3H, J=7.3 Hz, 14-Me), 2.29 (s, 6H, NMe<sub>2</sub>), 2.95 (s, 3H, 6-OMe), 5.06  
20 (d, 1H, J=11.0 Hz, H-3), 5.48 (dd, 1H, J=11.0, 1.8 Hz, H-13).

### Reference Example 3

#### Preparation of 2-aminoethylsulfonanilide

25 (1) To a solution of 8.0 g (64 mmol) of 2-aminoethylsulfonic acid (taurine) in 35 ml of distilled water were added 11.2 g (0.13 mol) of sodium bicarbonate and 19.8 g (0.12 mol) of carbobenzoxychloride under ice-

cooling, followed by stirring at room temperature for 4 hours. After the reaction, the reaction solution was washed with ether, the aqueous layer was made acidic with an aqueous hydrochloric acid solution, and the solution was concentrated under reduced pressure. The precipitated crystals were recrystallized from water/methanol (1:10) to give 15.5 g (yield: 86 %) of carbobenzoxytaurine.

10           (2) To a suspension of 15.0 g (53.3 mmol) of the compound obtained in the above (1) in 150 ml of benzene was added 15.0 g (72 mmol) of phosphorus pentachloride, followed by refluxing under heating for 25 minutes. After the reaction, the reaction solution was evaporated under reduced pressure, and 150 ml of benzene was added thereto, followed by removal of the insoluble matter by filtration. The filtrate was evaporated under reduced pressure to give 15.0 g of sulfonyl chloride.

20           (3) To a solution of 3.7 g (13.2 mmol) of the compound obtained in the above (2) in 100 ml of methylene chloride were added 1.8 ml (19.0 mmol) of aniline, 4.0 ml (51.6 mmol) of pyridine and 0.16 g (1.3 mmol) of 4-dimethylaminopyridine, followed by stirring overnight. After the reaction, the reaction solution was washed with 1N aqueous hydrochloric acid solution, the organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced

pressure. The residue was dissolved in 100 ml of methanol, and 0.5 g of palladium carbon was added thereto, followed by stirring under a hydrogen stream overnight. After the reaction, the catalyst in the  
5 reaction solution was removed by filtration, and the filtrate was concentrated under reduced pressure to give 1.7 g of 2-aminoethylsulfonanilide.

#### Example 34

10 11-[2-(N-Phenylsulfamoyl)ethyl]amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 33 using 0.38 g (0.50 mmol) of the compound obtained in  
15 Example 12(3), 1.0 g (5.0 mmol) of 2-aminoethylsulfonanilide described in Reference Example 3, there was obtained 0.34 g (yield: 85 %) of the title compound.  
FABMS m/z: 796 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ (ppm); 0.87 (t, 3H, J=7.3 Hz,  
20 14-Me), 2.26 (s, 6H, NMe<sub>2</sub>), 2.60 (s, 3H, 6-OMe), 4.23 (d, 1H, J=8.5 Hz, H-5).

#### Example 35

11-[2-[N-(4-Methoxyphenyl)sulfamoyl]ethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate  
25

Following the same procedure as in Example 33 using 0.50 g (0.57 mmol) of the compound obtained in

Example 10(4), 4-methoxyaniline in place of aniline and 1.30 g (5.7 mmol) of 4'-methoxy-2-aminoethyl-sulfonanilide prepared in the same manner as in Reference Example 3, there was obtained 0.06 g (yield: 11 %) of the title compound.

FABMS m/z: 947 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ (ppm); 0.83 (t, 3H, J=7.3 Hz, 14-Me), 2.29 (s, 6H, NMe<sub>2</sub>), 2.97 (s, 3H, 6-OMe), 3.79 (s, 3H, Ph-OMe), 5.06 (d, 1H, J=10.6 Hz, H-3), 5.48 (dd, 1H, J=11.0, 2.0 Hz, H-13), 6.87 (m, 2H), 7.29 (m, 2H).

#### Example 36

11-[2-[N-(3-Pyridyl)sulfamoyl]ethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 33 using 0.35 g (0.40 mmol) of the compound obtained in Example 10(4), 3-aminopyridine in place of aniline and 0.80 g (3.98 mmol) of 3-(2-aminoethyl)sulfonamidopyridine prepared in the same manner as in Reference Example 3, there was obtained 0.05 g (yield: 14 %) of the title compound.

FABMS m/z: 918 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ (ppm); 0.84 (t, 3H, J=7.3 Hz, 14-Me), 2.29 (s, 6H, NMe<sub>2</sub>), 2.99 (s, 3H, 6-OMe), 5.06 (d, 1H, J=11.0 Hz, H-3), 5.48 (dd, 1H, J=11.2, 2.0 Hz, H-13), 6.87 (m, 2H), 7.29 (m, 2H).

## Example 37

11-[2-[(4-Nitrophenylsulfonyl)amino]ethyl]-  
amino-11-deoxy-3-O-methylthiomethyl-5-O-desosaminyl-6-O-  
methylerythronolide A 11,12-cyclic carbamate

5 (1) Following the same procedures as in Example 1(1) and (2) using 5.06 g (7.69 mmol) of the compound obtained in Reference Example 1 and 2.3 g (10.4 mmol) of 4-nitrobenzenesulfonyl chloride, there was obtained 7.31 g of the 2'-O-acetyl compound.

10 (2) Following the same procedure as in Example 2 using 4.02 g (4.54 mmol) of the compound obtained in the above (1), there were obtained 1.33 g (yield: 35 %) of the same compound as obtained in Example 14 and 0.15 g (yield: 4 %) of the title compound.

15 IonSpray MS m/z: 903.4[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 2.28(s, 6H, NMe<sub>2</sub>), 2.29(s, 3H, SMe), 2.94(s, 3H, 6-OMe), 4.44(d, 1H, J=7.3Hz, H-1'), 4.68 and 4.92(each d, each 1H, J=11.0Hz, -OCH<sub>2</sub>SMe), 4.86(dd, 1H, J=11.0, 1.8Hz, H-13)

20 <sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 15.8(SMe), 40.2(NMe<sub>2</sub>), 50.2(6-OMe), 77.7(-OCH<sub>2</sub>SMe), 101.7(C1'), 158.0(11, 12-carbamate), 176.5(C1), 216.2(C9).

## Example 38

25 11-[2-(3-Pyridylsulfonylamino)ethyl]amino-  
3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methyl-  
erythronolide A 11,12-cyclic carbamate

(1) Following the same procedures as in



Example 1(1) and (2) using 1.00 g (1.52 mmol) of the compound obtained in Reference Example 1 and 0.60 g (3.4 mmol) of 3-pyridylsulfonyl chloride obtained in Reference Example 2, there was obtained 0.42 g of the  
 5 2'-O-acetyl compound.

(2) Following the same procedure as in Example 2 using 0.42 g (0.5 mmol) of the compound obtained in the above (1), there was obtained 0.21 g (yield: 17 %) of the title compound.

10 IonSpray MS m/z: 797.4[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 2.26(s, 6H, NMe<sub>2</sub>), 2.51(s, 3H, 6-OMe), 3.84(q, 1H, J=6.7Hz, H-2), 4.22(d, 1H, J=9.2Hz, H-5), 4.27(d, 1H, J=7.3Hz, H-1'), 4.86(dd, 1H, J=11.0, 2.4Hz, H-13), 6.12(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)

15 <sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 40.2(NMe<sub>2</sub>), 49.7(6-OMe), 103.9(C1'), 158.0(11,12-carbamate), 170.2(C1), 203.2(C3), 216.6(C9).

#### Example 39

20 11-[2-[(5-[2-(Methylthio)pyrimidin-4-yl]thiophene-2-sulfonylamino]ethylamino-11-deoxy-3-O-(2-pyridyl)-acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example  
 25 10(6) using 0.55 g (0.71 mmol) of the compound obtained in Example 10(5) and 0.26 g (0.85 mmol) of 5-[2-(methylthio)pyrimidin-4-yl]thiophene-2-sulfonyl chloride, there was obtained 0.50 g (yield: 67 %) of the title

compound.

FABMS m/z: 1047[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 2.30(s, 6H, NMe<sub>2</sub>), 2.60(s, 3H, SMe), 2.91(s, 3H, 6-OMe), 3.92 and 3.96(each d, each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 4.26(d, 1H, J=7.3Hz, H-1'), 4.98(dd, 1H, J=11.0, 1.8Hz, H-13), 5.00(d, 1H, J=11.6Hz, H-3), 6.24(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 14.1(SMe), 40.3(NMe<sub>2</sub>), 50.1(6-OMe), 103.5(C1'), 157.6(11,12-carbamate), 170.4(-COCH<sub>2</sub>[2-Pyridine]), 175.2(C1), 216.0(C9).

#### Example 40

11-[2-(6-Chloroimidazo[2,1-b]thiazole-5-sulfonylamino)ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.62 g (0.80 mmol) of the compound obtained in Example 10(5) and 0.26 g (1.0 mmol) of 6-chloroimidazo[2,1-b]thiazole-5-sulfonyl chloride, there was obtained 0.16 g (yield: 19 %) of the title compound.  
 FABMS m/z: 997[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 2.30(s, 6H, NMe<sub>2</sub>), 2.85(s, 3H, 6-OMe), 3.92 and 3.96(each d, each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 4.06(d, 1H, J=7.3Hz, H-1'), 4.99(d, 1H, J=11.0Hz, H-3), 5.00(dd, 1H, J=11.0, 2.4Hz, H-13), 6.65(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 40.3(NMe<sub>2</sub>), 50.0(6-OMe),

103.5(C1'), 157.8(11,12-carbamate), 170.4(-COCH<sub>2</sub>[2-Pyridine]), 175.3(C1), 215.9(C9).

#### Example 41

5            11-[2-(5-Chloro-3-methylbenzo[b]thiophene-2-sulfonylamino)ethylamino-11-deoxy-3-O-(2-pyridyl)-acetyl-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example

10 10(6) using 0.51 g (0.65 mmol) of the compound obtained in Example 10(5) and 0.22 g (0.78 mmol) of 5-chloro-3-methylbenzo[b]thiophene-2-sulfonyl chloride, there was obtained 0.45 g (yield: 67%) of the title compound.

FABMS m/z:1021[M+H]<sup>+</sup>

15    <sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 2.29(s, 6H, NMe<sub>2</sub>), 2.66(s, 3H, benzothiophene-Me), 2.88(s, 3H, 6-OMe), 3.93 and 3.96(each d, each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 4.05(d, 1H, J=6.7Hz, H-1'), 4.97-5.00(m, 1H, H-13), 5.01(d, 1H, J=11.6Hz, H-3), 6.48(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)

20    <sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 12.2(benzothiophene-Me), 40.3(NMe<sub>2</sub>), 50.1(6-OMe), 103.5(C1'), 157.9(11,12-carbamate), 170.4(-COCH<sub>2</sub>[2-Pyridine]), 175.2(C1), 215.9(C9).

#### 25 Example 42

11-[2-(4-Methylsulfonylbenzenesulfonylamino)-ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic

carbamate

Following the same procedure as in Example 10(6) using 0.53 g (0.68 mmol) of the compound obtained in Example 10(5) and 0.23 g (0.90 mmol) of 4-methylsulfonylbenzenesulfonyl chloride, there was obtained 0.36 g (yield: 53 %) of the title compound.

FABMS  $m/z$ : 995  $[M+H]^+$

$^1H$ -NMR (500MHz,  $CDCl_3$ )  $\delta$  (ppm) 2.29 (s, 6H,  $NMe_2$ ), 2.87 (s, 3H, 6-OMe), 3.08 (s, 3H,  $SO_2Me$ ), 3.93 and 3.96 (each d, each 1H,  $J=15.9Hz$ ,  $-CH_2[2-Pyridine]$ ), 4.07 (d, 1H,  $J=7.3Hz$ , H-1'), 4.92 (dd, 1H,  $J=11.0, 1.8Hz$ , H-13), 5.00 (d, 1H,  $J=11.0Hz$ , H-3), 6.30 (brs, 1H,  $-NH-SO_2-$ )

$^{13}C$ -NMR (125MHz,  $CDCl_3$ )  $\delta$  (ppm) 40.3 ( $NMe_2$ ), 44.2 ( $SO_2Me$ ), 50.1 (6-OMe), 103.5 (C1'), 157.8 (11,12-carbamate), 170.6 ( $-COCH_2[2-Pyridine]$ ), 175.5 (C1), 216.0 (C9).

## Example 43

11-[2-(2-Thiophenesulfonylamino)ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.52 g (0.66 mmol) of the compound obtained in Example 10(5) and 0.15 g (0.82 mmol) of 2-thiophenesulfonyl chloride, there was obtained 0.46 g (yield: 75 %) of the title compound.

FABMS  $m/z$ : 923  $[M+H]^+$

$^1H$ -NMR (500MHz,  $CDCl_3$ )  $\delta$  (ppm) 2.29 (s, 6H,  $NMe_2$ ), 2.88 (s, 3H, 6-OMe), 3.92 and 3.96 (each d, each 1H,  $J=15.9Hz$ ,

-CH<sub>2</sub>[2-Pyridine]], 4.06(d, 1H, J=7.3Hz, H-1'),  
 4.98(dd, 1H, J=11.0, 1.8Hz, H-13), 5.01(d, 1H,  
 J=11.0Hz, H-3), 6.10(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm)40.3(NMe<sub>2</sub>), 50.1(6-OMe),  
 5 103.5(C1'), 157.7(11,12-carbamate), 170.5(-COCH<sub>2</sub>[2-  
 Pyridine]), 175.1(C1), 215.9(C9).

## Example 44

11-[2-[5-(Isoxazol-3-yl)thiophene-2-  
 10 sulfonylamino]ethylamino-11-deoxy-3-O-(2-  
pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A  
11,12-cyclic carbamate

Following the same procedure as in Example  
 10(6) using 0.58 g (0.75 mmol) of the compound obtained  
 15 in Example 10(5) and 0.22 g (0.88 mmol) of 5-(isoxazol-  
 3-yl)thiophene-2-sulfonyl chloride, there was obtained  
 0.36 g (yield: 48 %) of the title compound.

FABMS m/z:990[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm)2.29(s, 6H, NMe<sub>2</sub>), 2.93(s,  
 20 3H, 6-OMe), 3.91 and 3.95(each d, each 1H, J=15.9Hz,  
 -CH<sub>2</sub>[2-Pyridine]), 4.05(d, 1H, J=7.3Hz, H-1'),  
 4.95(dd, 1H, J=11.0, 2.2Hz, H-13), 5.01(d, 1H,  
 J=10.9Hz, H-3), 6.30(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm)40.3(NMe<sub>2</sub>), 50.2(6-OMe),  
 25 103.5(C1'), 157.9(11,12-carbamate), 170.4(-COCH<sub>2</sub>[2-  
 Pyridine]), 175.2(C1), 216.0(C9).

## Example 45

11-[2-(Benzofurazan-4-sulfonylamino)ethyl]-  
amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-  
O-methylerythronolide A 11,12-cyclic carbamate

5           Following the same procedure as in Example  
 10(6) using 0.58 g (0.75 mmol) of the compound obtained  
 in Example 10(5) and 0.19 g (0.87 mmol) of benzofurazan-  
 4-sulfonyl chloride, there was obtained 0.47 g (yield:  
 66 %) of the title compound.

10   FABMS m/z:990[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm)2.29(s, 6H, NMe<sub>2</sub>), 3.02(s,  
 3H, 6-OMe), 3.93 and 3.96(each d, each 1H, J=15.9Hz,  
 -CH<sub>2</sub>[2-Pyridine]), 4.08(d, 1H, J=7.3Hz, H-1'),  
 5.02(dd, 1H, J=11.0Hz, H-3), 5.11(dd, 1H, J=11.0,  
 15   1.8Hz, H-13), 6.48(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm)40.3(NMe<sub>2</sub>), 50.5(6-OMe),  
 103.5(C1'), 157.1(11,12-carbamate), 170.4(-COCH<sub>2</sub>[2-  
 Pyridine]), 175.3(C1), 215.8(C9).

## 20   Example 46

11-[2-(2-Methylsulfonylbenzenesulfonylamino)-  
ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-  
desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
carbamate

25           Following the same procedure as in Example  
 10(6) using 0.52 g (0.67 mmol) of the compound obtained  
 in Example 10(5) and 0.21 g (0.82 mmol) of 2-  
 methylsulfonylbenzenesulfonyl chloride, there was

obtained 0.46 g (yield: 66 %) of the title compound.

FABMS m/z: 995 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500MHz, CDCl<sub>3</sub>) δ (ppm) 2.29 (s, 6H, NMe<sub>2</sub>), 2.92 (s, 3H, 6-OMe), 3.92 and 3.96 (each d, each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 3.39 (s, 3H, SO<sub>2</sub>Me), 4.05 (d, 1H, J=7.3Hz, H-1'), 5.00 (d, 1H, J=11.0Hz, H-3), 5.04 (dd, 1H, J=11.0, 1.8Hz, H-13), 6.40 (brt, 1H, J=5.5Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125MHz, CDCl<sub>3</sub>) δ (ppm) 40.3 (NMe<sub>2</sub>), 50.1 (6-OMe), 103.6 (C1'), 157.2 (11,12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 174.0 (C1), 215.6 (C9).

#### Example 47

11-[2-(4-Nitrobenzenesulfonylamino)butylamino]-3,11-dideoxy-3-oxo-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate

(1) Following the same procedure as in Example 10(5) using 3.0 g (4.25 mmol) of the compound obtained in Example 12(3) and 4.3 ml (43 mmol) of 1,4-diaminobutane, there was obtained 3.2 g of 11-(3-aminobutyl)amino-3,11-dideoxy-3-oxo-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate.

(2) Following the same procedure as in Example 10(6) using 0.61 g (0.88 mmol) of the compound obtained in the above (1) and 0.26 g (1.2 mmol) of 4-nitrobenzenesulfonyl chloride, there was obtained 0.45 g (yield: 58 %) of the title compound.

FABMS m/z: 869 [M+H]<sup>+</sup>

- <sup>1</sup>H-NMR (500MHz, CDCl<sub>3</sub>) δ (ppm) 2.27 (s, 6H, NMe<sub>2</sub>), 2.60 (s, 3H, 6-OMe), 3.87 (q, 1H, J=6.7Hz, H-2), 4.26 (d, 1H, J=8.6Hz, H-5), 4.29 (d, 1H, J=7.3Hz, H-1'), 4.88 (dd, 1H, J=11.0, 1.8Hz, H-13), 5.65 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)
- 5    <sup>13</sup>C-NMR (125MHz, CDCl<sub>3</sub>) δ (ppm) 40.2 (NMe<sub>2</sub>), 49.9 (6-OMe), 104.0 (C1'), 157.3 (11,12-carbamate), 170.1 (C1), 203.4 (C3), 216.9 (C9).

## Example 48

- 10                    11-[2-(4-Nitrobenzenesulfonylamino)butyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

- (1) Following the same procedure as in Example 10(5) using 5.0 g (6.1 mmol) of the compound obtained in Example 10(4) and 6.1 ml (61 mmol) of 1,4-diaminopropane, there was obtained 4.9 g of 11-(3-aminobutyl)amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate.

- (2) Following the same procedure as in Example 20 10(6) using 0.45 g (0.56 mmol) of the compound obtained in the above (1) and 0.14 g (0.63 mmol) of 4-nitrobenzenesulfonyl chloride, there was obtained 0.34 g (yield: 62 %) of the title compound.

FABMS m/z: 990 [M+H]<sup>+</sup>

- 25    <sup>1</sup>H-NMR (500MHz, CDCl<sub>3</sub>) δ (ppm) 2.30 (s, 6H, NMe<sub>2</sub>), 3.00 (s, 3H, 6-OMe), 3.93 and 3.99 (each d, each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 4.08 (d, 1H, J=7.3Hz, H-1'), 4.97 (dd, 1H, J=11.0, 2.1Hz, H-13), 5.06 (d, 1H,



J=11.2Hz, H-3).

Example 49

11-[2-(2-Acetamido-4-methyl-5-  
 5 thiazolesulfonylamino)butyllamino-11-deoxy-3-O-(2-  
pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A  
11,12-cyclic carbamate

Following the same procedure as in Example  
 10(6) using 0.43 g (0.53 mmol) of the compound obtained  
 10 in Example 48(1) and 0.16 g (0.63 mmol) of 2-acetamido-  
 4-methyl-5-thiazolesulfonyl chloride, there was obtained  
 0.38 g (yield: 70 %) of the title compound.

FABMS m/z:1023[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 2.23(s, 3H, COCH<sub>3</sub>), 2.29(s,  
 15 6H, NMe<sub>2</sub>), 2.98(s, 3H, 6-OMe), 3.93 and 3.98(each d,  
 each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 4.08(d, 1H,  
 J=7.3Hz, H-1'), 4.98(dd, 1H, J=11.0, 2.1Hz, H-13),  
 5.02(d, 1H, J=11.3Hz, H-3), 5.63(brt, 1H, J=5.8Hz,  
 -NH<sub>2</sub>SO<sub>2</sub>-)

20 <sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 22.9(COCH<sub>3</sub>), 40.3(NMe<sub>2</sub>),  
 50.2(6-OMe), 103.6(C1'), 157.5(11, 12-carbamate),  
 168.2(COCH<sub>3</sub>), 170.7(-COCH<sub>2</sub>[2-Pyridine]), 174.5(C1),  
 216.4(C9).

25 Example 50

11-[2-(3-Pyridylsulfonylamino)butyllamino-11-  
deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-  
methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.58 g (0.72 mmol) of the compound obtained in Example 48(1) and 0.38 g (2.14 mmol) of 3-pyridylsulfonyl chloride obtained in Reference Example 2, there was obtained 0.51 g (yield: 75 %) of the title compound.

FABMS m/z: 946 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500MHz, CDCl<sub>3</sub>) δ (ppm) 2.30 (s, 6H, NMe<sub>2</sub>), 3.04 (s, 3H, 6-OMe), 3.93 and 3.97 (each d, each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 4.08 (d, 1H, J=7.3Hz, H-1'), 4.98 (dd, 1H, J=11.0, 1.8Hz, H-13), 5.06 (d, 1H, J=11.0Hz, H-3), 5.77 (brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)

<sup>13</sup>C-NMR (125MHz, CDCl<sub>3</sub>) δ (ppm) 40.3 (NMe<sub>2</sub>), 50.4 (6-OMe), 103.6 (C1'), 157.4 (11, 12-carbamate), 170.4 (-COCH<sub>2</sub>[2-Pyridine]), 174.8 (C1), 216.9 (C9).

#### Example 51

11-[2-(3-Pyridylsulfonylamino)butyl]amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythronolide A  
11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.78 g (1.15 mmol) of the compound obtained in Example 47(1) and 0.61 g (3.43 mmol) of 3-pyridylsulfonyl chloride obtained in Reference Example 2, there was obtained 0.39 g (yield: 41 %) of the title compound.

FABMS m/z: 825 [M+H]<sup>+</sup>

<sup>1</sup>H-NMR (500MHz, CDCl<sub>3</sub>) δ (ppm) 2.27 (s, 6H, NMe<sub>2</sub>), 2.63 (s,

3H, 6-OMe), 3.87(q, 1H, J=6.7Hz, H-2), 4.27(d, 1H, J=10.4Hz, H-5), 4.29(d, 1H, J=7.3Hz, H-1'), 4.90(dd, 1H, J=11.0, 1.8Hz, H-13), 5.68(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-)  
<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 40.2(NMe<sub>2</sub>), 50.0(6-OMe),  
 5 103.9(C1'), 157.3(11,12-carbamate), 170.1(C1),  
 203.4(C3), 216.9(C9).

## Example 52

11-[2-(2-Dibenzofuransulfonylamino)ethyl]amino-3,11-dideoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.33 g (0.42 mmol) of the compound obtained in Example 10(5) and 0.13 g (0.49 mmol) of 2-dibenzofuransulfonyl chloride, there was obtained 0.24 g (yield: 57 %) of the title compound.

FABMS m/z: 1007[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(300MHz, CDCl<sub>3</sub>) δ(ppm) 2.29(s, 6H, NMe<sub>2</sub>), 2.88(s, 3H, 6-OMe), 3.97 and 3.91(each d, each 1H, J=15.9Hz, -CH<sub>2</sub>[2-Pyridine]), 4.05(d, 1H, J=7.3Hz, H-1'), 4.96(dd, 1H, J=11.0, 2.0Hz, H-13), 5.01(d, 1H, J=11.2Hz, H-3), 6.00(brt, 1H, -NH<sub>2</sub>SO<sub>2</sub>-).

## Example 53

11-[2-(2-Methylsulfonylbenzenesulfonylamino)-ethyl]amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example

12(5) using 0.51 g (0.73 mmol) of the compound obtained in Example 12(4) and 0.22 g (0.86 mmol) of 2-methylsulfonylbenzenesulfonyl chloride, there was obtained 0.46 g (yield: 72 %) of the title compound.

5 IonSprayMS m/z:874.2[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(300MHz, CDCl<sub>3</sub>) δ(ppm)2.26(s, 6H, NMe<sub>2</sub>), 2.55(s, 3H, 6-OMe), 3.40(s, 3H, SO<sub>2</sub>Me), 4.96(dd, 1H, J=10.6, 2.5Hz, H-13), 6.37(brt, 1H, J=6.2Hz, -NH<sub>2</sub>SO<sub>2</sub>-)

10 Example 54

11-[2-[2-(Pyrid-2-yl)thiophene-5-sulfonylamino]ethylamino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

15 Following the same procedure as in Example 12(5) using 0.51 g (0.73 mmol) of the compound obtained in Example 12(4) and 0.24 g (0.92 mmol) of 2-(pyrid-2-yl)thiophene-5-sulfonyl chloride, there was obtained 0.33 g (yield: 48 %) of the title compound.

20 IonSprayMS m/z:879.3[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(300MHz, CDCl<sub>3</sub>) δ(ppm)2.26(s, 6H, NMe<sub>2</sub>), 2.58(s, 3H, 6-OMe), 4.22(d, 1H, J=8.7Hz, H-5), 4.27(d, 1H, J=7.3Hz, H-1'), 4.92(dd, 1H, J=10.6, 2.3Hz, H-13), 6.00(brs, 1H, -NH<sub>2</sub>SO<sub>2</sub>-).

25

Example 55

11-[2-(2-Methylsulfonylbenzenesulfonylamino)-ethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-

desosaminylerythronolide A 11,12-cyclic carbamate

(1) Following the same procedure as in Example 10(5) using 2.47 g of 2'-O-acetyl-10,11-anhydro-12-O-imidazolylcarbonyl-3-O-(2-pyridyl)acetyl-5-O-desosaminylerythronolide A, there was obtained 2.29 g of 11-(2-aminoethyl)amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminylerythronolide A 11,12-cyclic carbamate.

(2) Following the same procedure as in Example 10(6) using 0.52 g (0.68 mmol) of the compound obtained in the above (1) and 0.21 g (0.82 mmol) of 2-methylsulfonylbenzenesulfonyl chloride, there was obtained 0.38 g (yield: 57 %) of the title compound.

FABMS m/z: 981[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 2.33(s, 6H, NMe<sub>2</sub>), 3.40(s, 3H, SO<sub>2</sub>Me), 3.91 and 3.95(each d, each 1H, J=15.3Hz, -CH<sub>2</sub>[2-Pyridine]), 4.57(d, 1H, J=7.3Hz, H-1'), 4.99 and 5.48(each brs, each 1H, H-3 and H-13), 6.25(brs, 1H, -NH-SO<sub>2</sub>-)

<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 40.5(NMe<sub>2</sub>), 44.3(SO<sub>2</sub>Me), 104.5(C1'), 156.5(11, 12-carbamate), 170.0(-COCH<sub>2</sub>[2-Pyridine]), 174.6(C1), 214.5(C9).

## Example 56

11-[2-[(2-Acetamido-4-methyl-5-thiazolesulfonyl)amino]ethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

(1) Following the same procedure as in Example

10(5) using 5.0 g (6.1 mmol) of the compound obtained in Example 10(4) and 4.5 g (61 mmol) of 1,3-diaminopropane, there was obtained 4.5 g of 11-(3-aminopropyl)amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate.

(2) Following the same procedure as in Example 10(6) using 0.50 g (0.63 mmol) of the compound obtained in the above (1) and 0.24 g (0.95 mmol) of 2-acetamido-4-methyl-5-thiazolesulfonyl chloride, there was obtained 0.47 g (yield: 74 %) of the title compound.

FABMS m/z:1009[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm) 0.76(t, 3H, J=7.3Hz, 14-Me), 2.29(s, 6H, NMe<sub>2</sub>), 2.93(s, 3H, 6-OMe), 4.07(d, 1H, J=7.3Hz, H-1'), 4.92(dd, 1H, J=11.0, 2.1Hz, H-13), 5.01(d, 1H, J=11.0Hz, H-3).

<sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm) 40.3(NMe<sub>2</sub>), 50.0(6-OMe), 103.7(C1'), 157.9(11,12-carbamate), 174.5(C1), 216.0(C9).

## Example 57

11-[3-(4-Nitrobenzenesulfonyl)aminopropyl]-amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminy-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.50 g (0.63 mmol) of the compound obtained in Example 56(1) and 0.21 g (0.95 mmol) of 4-nitrobenzenesulfonyl chloride, there was obtained 0.45 g (yield: 21 %) of the title compound.

FABMS m/z:976[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm)2.29(s, 6H, NMe<sub>2</sub>), 2.98(s, 3H, 6-OMe), 4.06(d, 1H, J=7.3Hz, H-1'), 4.83(dd, 1H, J=11.0, 1.8Hz, H-13), 5.02(d, 1H, J=11.0Hz, H-3).

5 <sup>13</sup>C-NMR(125MHz, CDCl<sub>3</sub>) δ(ppm)40.3(NMe<sub>2</sub>), 50.0(6-OMe), 103.6(C1'), 158.1(11,12-carbamate), 174.7(C1), 216.1(C9).

#### Example 58

10 11-[3-(3-Pyridylsulfonyl)aminopropyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.50 g (0.63 mmol) of the compound obtained in Example 56(1) and 0.40 g (2.3 mmol) of pyridine-3-sulfonyl chloride obtained in Reference Example 2, there was obtained 0.24 g (yield: 21 %) of the title compound.

FABMS m/z:932[M+H]<sup>+</sup>

20 <sup>1</sup>H-NMR(500MHz, CDCl<sub>3</sub>) δ(ppm)2.29(s, 6H, NMe<sub>2</sub>), 2.99(s, 3H, 6-OMe), 4.06(d, 1H, J=7.3Hz, H-1'), 4.86(dd, 1H, J=11.0, 2.4Hz, H-13), 5.02(d, 1H, J=11.6Hz, H-3), 5.89(brs, 1H, -NHSO<sub>2</sub>-).

#### Example 59

25 11-[3-[(2-Acetamido-4-methyl-5-thiazolesulfonyl)aminopropyl]amino-3,11-dideoxy-3-oxo-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

(1) Following the same procedure as in Example 10(5) using 2.0 g (2.8 mmol) of the compound obtained in Example 12(3) and 2.1 g (28 mmol) of 1,3-diaminopropane, there was obtained 2.0 g of 11-(3-aminopropyl)amino-  
5 3,11-dideoxy-3-oxo-5-O-desosaminy1-6-O-methylerythronolide A 11,12-cyclic carbamate.

(2) Following the same procedure as in Example 10(6) using 0.40 g (0.56 mmol) of the compound obtained in the above (1) and 0.22 g (0.84 mmol) of 2-acetamido-  
10 4-methyl-5-thiazolesulfonyl chloride, there was obtained 0.31 g (yield: 62 %) of the title compound.

FABMS  $m/z$ : 888  $[M+H]^+$

$^1\text{H-NMR}$  (500MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm) 2.27 (s, 6H,  $\text{NMe}_2$ ), 2.62 (s, 3H, 6-OMe), 4.23 (d, 1H,  $J=9.2\text{Hz}$ , H-5), 4.28 (d, 1H,  $J=7.3\text{Hz}$ , H-1'), 4.84 (dd,  $J=11.0$ , 2.4Hz, H-13),  
15 5.89 (brs, 1H,  $-\text{NH}\text{SO}_2-$ )

$^{13}\text{C-NMR}$  (125MHz,  $\text{CDCl}_3$ )  $\delta$  (ppm) 40.2 ( $\text{NMe}_2$ ), 49.9 (6-OMe), 104.0 ( $\text{C1}'$ ), 158.0 (11, 12-carbamate), 169.8 ( $\text{C1}$ ), 203.5 ( $\text{C3}$ ), 216.4 ( $\text{C9}$ ).

20

#### Example 60

11-[3-(3-Pyridylsulfonyl)aminopropyl]amino-3,11-dideoxy-3-oxo-5-O-desosaminy1-6-O-methylerythronolide A 11,12-cyclic carbamate

25 Following the same procedure as in Example 10(6) using 0.39 g (0.58 mmol) of the compound obtained in Example 59(1) and 0.22 g (0.84 mmol) of 2-acetamido-4-methyl-5-thiazolesulfonyl chloride, there was obtained



0.31 g (yield: 51 %) of the title compound.

FABMS  $m/z$ : 811  $[M+H]^+$

$^1H$ -NMR (300MHz,  $CDCl_3$ )  $\delta$  (ppm) 0.76 (t, 1H,  $J=7.5$ Hz, 14-Me),  
2.27 (s, 6H,  $NMe_2$ ), 2.60 (s, 3H, 6-OMe), 4.23 (d, 1H,  
5  $J=8.6$ Hz, H-5), 4.28 (d, 1H,  $J=7.3$ Hz, H-1'), 4.78 (dd,  
 $J=11.0$ , 2.4Hz, H-13), 5.82 (brs, 1H,  $-NH_2SO_2^-$ )

#### Example 61

11-[2-[N-(2-Methylsulfonylbenzene)sulfonyl-N-  
10 methylaminoethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-  
5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
carbamate

(1) 8.0 g (9.14 mmol) of 11-[2-(N-methyl-N-  
benzylamino)ethyl]amino-11-deoxy-3-O-(2-pyridyl)acetyl-  
15 5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
carbamate was hydrogenated in methanol using palladium-  
carbon in an ordinary manner. The reaction solution was  
filtered, and the solvent was evaporated under reduced  
pressure to give 5.0 g (yield: 69 %) of the debenzylated  
20 compound.

(2) Following the same procedure as in Example  
10(6) using 0.40 g (0.51 mmol) of the compound obtained  
in the above (1) and 0.16 g (0.61 mmol) of 2-  
methylsulfonylbenzenesulfonyl chloride, there was  
25 obtained 0.39 g (yield: 76 %) of the title compound.

SIMS  $m/z$ : 1009  $[M+H]^+$

$^1H$ -NMR (300MHz,  $CDCl_3$ )  $\delta$  (ppm) 0.82 (t, 3H,  $J=7.3$ Hz, 14-Me),  
2.29 (s, 6H,  $NMe_2$ ), 2.99 (s, 3H, 6-OMe), 4.06 (d, 1H,

J=7.3Hz, H-1'), 5.03-5.11(m, 2H, H-3 and H-13).

Example 62

5 11-[2-[N-(6-Chloroimidazo[2,1,6]thiazole-5-sulfonyl)-N-methylaminoethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.40 g (0.51 mmol) of the compound obtained in Example 61 and 0.16 g (0.61 mmol) of 6-chloroimidazo-[2,1,6]thiazole-5-sulfonyl chloride, there was obtained 0.38 g (yield: 74 %) of the title compound.

SIMS m/z:1011[M+H]<sup>+</sup>

<sup>1</sup>H-NMR(300MHz, CDCl<sub>3</sub>) δ(ppm)0.81(t, 3H, J=7.3Hz, 14-Me),  
15 2.30(s, 6H, NMe<sub>2</sub>), 3.03(s, 3H, 6-OMe), 4.06(d, 1H, J=7.3Hz, H-1'), 5.03-5.11(m, 2H, H-3 and H-13).

Example 63

20 11-[2-[N-(3-Pyridylsulfonyl)-N-methylaminoethylamino-11-deoxy-3-O-(2-pyridyl)acetyl-5-O-desosaminyl-6-O-methylerythronolide A 11,12-cyclic carbamate

Following the same procedure as in Example 10(6) using 0.40 g (0.51 mmol) of the compound obtained in Example 61 and 0.18 g (1.02 mmol) of pyridine-3-sulfonyl chloride, there was obtained 0.25 g (yield: 53 %) of the title compound.

SIMS m/z:932[M+H]<sup>+</sup>

<sup>1</sup>H-NMR (300MHz, CDCl<sub>3</sub>) δ (ppm) 0.83 (t, 3H, J=7.3Hz, 14-Me),  
2.30 (s, 6H, NMe<sub>2</sub>), 2.97 (s, 3H, 6-OMe), 4.07 (d, 1H,  
J=7.3Hz, H-1'), 5.04 (dd, 1H, J=11.0, 2.3Hz, H-13),  
5.05 (d, 1H, J=11.2Hz, H-3).

5

## Example 64

11-[2-[(2-Nitrophenyl)sulfonylamino]ethyl]-  
amino-11-deoxy-3-O-[(3-pyridylmethyl)aminolcarbonyl-5-O-  
desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
10 carbamate

To a solution of 2.2 g (2.5 mmol) of the  
compound obtained in Example 3(1) in 10 ml of pyridine  
was added dropwise a solution of 0.74 g (2.5 mmol) of  
triphosgene in 10 ml of methylene chloride under ice-  
15 cooling, followed by stirring for 30 minutes. To the  
mixture was added 1.25 ml (12.3 mmol) of 3-  
(aminomethyl)pyridine, followed by stirring for 1.5  
hours. After completion of the reaction, the same  
working-up and removal of the acetyl group at the 2'-  
20 position as in Example 1(3) gave 1.6 g of the title  
compound.

FABMS m/z: 977 [M+H]<sup>+</sup>

## Example 65

25 11-[2-[(2-Nitrophenyl)sulfonylamino]-  
ethylamino-11-deoxy-3-O-(3-pyridyloxy)carbonyl-5-O-  
desosaminyl-6-O-methylerythronolide A 11,12-cyclic  
carbamate

To a solution of 2.2 g (2.5 mmol) of the compound obtained in Example 3(1) in 10 ml of pyridine was added dropwise a solution of 0.74 g (2.5 mmol) of triphosgene in 10 ml of methylene chloride under ice-cooling, followed by stirring for 30 minutes. To the mixture was added 1.18 g (12.4 mmol) of 3-hydroxypyridine, followed by stirring for 1.5 hours. After completion of the reaction, the same working-up and removal of the acetyl group at the 2'-position as in Example 1(3) gave 1.2 g of the title compound.

FABMS m/z: 964 [M+H]<sup>+</sup>

#### Test Example

The in vitro antibacterial activity of the compound obtained in Example 13 as an example of the compound of the present invention against various experimental bacteria was measured using sensitive disc media (produced by Eiken Chemical Co.) according to the MIC measuring method specified by the Japan Society of Chemotherapy. The results are expressed as MIC value (Minimum Inhibitory Concentration,  $\mu\text{g/ml}$ ), and shown in Table 1.

[Table 1]

In Vitro Antibacterial Activity: MIC ( $\mu\text{g/ml}$ )

Compound Microorganism	Compound of Example 13
<i>S. aureus</i> 209P-JC	0.10
<i>S. aureus</i> Smith	0.20
<i>S. epidermidis</i> IID 866	0.025
<i>E. faecalis</i> CSJ 1212	0.10
<i>S. pneumoniae</i> BM 225	0.78
<i>S. pneumoniae</i> BM 205	1.56


## INDUSTRIAL APPLICABILITY

5           The compounds of the present invention have a strong antibacterial activity against not only erythromycin-sensitive bacteria but also erythromycin-resistant bacteria. Therefore, the compounds of the present invention are useful as antibacterial agents for


10 the treatment of bacterially infectious diseases in human beings and animals (including farm animals).

## CLAIMS

1. An erythromycin A derivative represented by the formula:




- 5 wherein n is an integer of 1 to 7,  
 $R^1$  is a group represented by the formula:




- wherein  $R^7$  is a hydrogen atom, an alkyl group having 1 to 6 carbon atoms, a phenyl group, a phenyl group substituted by a nitro group or an alkoxy group having 1 to 3 carbon atoms, a pyridyl group, a pyridyl group substituted by 1 or 2 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms; a halogen atom; an alkoxy group having 1 to 3 carbon atoms; a nitro group, an amino group; a cyano group and an amino group substituted by an alkyl group having 1 to 6 carbon atoms, a quinolyl group, or a quinolyl group substituted by 1 or 2 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms; a halogen atom; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group and an amino group substituted by an alkyl group having 1 to

6 carbon atoms,  $R^8$  is a hydrogen atom or an alkyl group having 1 to 6 carbon atoms, or a group represented by the formula:




- 5 wherein  $R^9$  is an alkyl group having 1 to 6 carbon atoms, a dibenzofuranyl group, a thienyl group, a thienyl group substituted by a group selected from the group consisting of a pyridyl group; an isoxazolyl group; a pyrimidinyl group and a pyrimidinyl group substituted by an alkoxy
- 10 group having 1 to 6 carbon atoms or an alkylthio group having 1 to 6 carbon atoms, an isoxazolyl group, an isoxazolyl group substituted by 1 or 2 alkyl groups having 1 to 6 carbon atoms, an imidazolyl group, an imidazolyl group substituted by 1 to 3 alkyl groups having 1 to 6
- 15 carbon atoms, a benzothienyl group, a benzothienyl group substituted by 1 to 5 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms and a halogen atom, a thiazolyl group, a thiazolyl group substituted by 1 or 2 members selected from the group
- 20 consisting of an alkyl group having 1 to 6 carbon atoms; an amino group and an acetamino group, an imidazo[2,1-b]thiazolyl group, an imidazo[2,1-b]thiazolyl group substituted by 1 to 3 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms
- 25 and a halogen atom, a phenylalkyl group having 7 to 10 carbon atoms, a quinolyl, a pyridyl, a naphthyl group, a naphthylalkyl group having 11 to 15 carbon atoms, a dimethylaminonaphthyl group, a group represented by the

formula:



wherein X is -O- or -S-,

a group represented by the formula:



5

a phenyl group, a phenyl group substituted by 1 to 5 members selected from the group consisting of a hydroxyl group; a methylsulfonyl group; an alkyl group having 1 to 6 carbon atoms; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group; a dimethylamino group; an acetylamino group; a pyridyl group; a trifluoromethyl group; a trifluoromethoxy group and a halogen atom, a pyridyl group, a pyridyl group substituted by 1 or 2 members selected from the group consisting of a hydroxyl group; an alkyl group having 1 to 6 carbon atoms; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano group; a dimethylamino group; an acetylamino group; a pyridyl group; a trifluoromethyl group; a trifluoromethoxy group and a halogen atom, a quinolyl group, or a quinolyl group substituted by 1 or 2 members selected from the group consisting of a hydroxyl group; an alkyl group having 1 to 6 carbon atoms; an alkoxy group having 1 to 3 carbon atoms; a nitro group; an amino group; a cyano

10

15


20




group; a dimethylamino group; an acetylamino group; a pyridyl group; a trifluoromethyl group; a trifluoromethoxy group and a halogen atom, and R<sup>10</sup> is a hydrogen atom or an alkyl group having 1 to 6 carbon atoms,

R<sup>2</sup> is a hydrogen atom, an alkyl group having 1 to 6 carbon atoms or a cinnamyl group,


R<sup>3</sup> is a group represented by the formula:



10 a group represented by the formula:




a group represented by the formula:



a group represented by the formula:

15  $-\text{O}-\text{R}^{11}$

or a group represented by the formula:



wherein R<sup>11</sup> is a pyridylmethyl group, a methylthiomethyl group, a quinolyl group, a phenyl group, a phenyl group substituted by 1 to 5 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms; a nitro group; an alkoxy group having 1 to 3 carbon atoms and a halogen atom, a pyridyl group, or a pyridyl group substituted by 1 or 2 members selected from the group consisting of an alkyl group having 1 to 6 carbon atoms; a nitro group; an alkoxy group having 1 to 3 carbon atoms and a halogen atom,

25 R<sup>4</sup> is a hydrogen atom, or R<sup>3</sup> and R<sup>4</sup> together form an oxo

group, and

R<sup>5</sup> and R<sup>6</sup> are the same or different, and are each a hydrogen atom or an alkyl group having 1 to 6 carbon atoms, or a pharmaceutically acceptable salt thereof.

- 5 2. A pharmaceutical composition comprising an effective amount of the erythromycin A derivative or the pharmaceutically acceptable salt thereof according to Claim 1.
3. An antibacterial preparation comprising the  
10 erythromycin A derivative or the pharmaceutically acceptable salt thereof according to Claim 1 as an effective component.
4. A method for the treatment of a bacterially  
15 pharmaceutically effective amount of the erythromycin A derivative or the pharmaceutically acceptable salt thereof according to Claim 1 to a patient.
5. Use of the erythromycin A derivative or the  
20 pharmaceutically acceptable salt thereof according to Claim 1 for the treatment of a bacterially infectious disease.

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP 98/04876

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07H17/08 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 596 802 A (ROUSSEL UCLAF) 11 May 1994 see the whole document ---	1-5
A	WO 97 31929 A (ROUSSEL UCLAF ; AGOURIDAS CONSTANTIN (FR); CHANTOT JEAN FRANCOIS (F) 4 September 1997 see the whole document -----	1-5

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"S" document member of the same patent family

Date of the actual completion of the international search

1 February 1999

Date of mailing of the international search report

12/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel.: (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Scott, J

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP 98/04876

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 4  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim 4  
is directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

information on patent family members

Intern. Appl. Application No.

PCT/JP 98/04876

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0596802 A	11-05-1994	FR 2697523 A	06-05-1994
		FR 2707088 A	06-01-1995
		AT 135706 T	15-04-1996
		AU 670329 B	11-07-1996
		AU 5047793 A	19-05-1994
		BR 9304481 A	21-06-1994
		CA 2102457 A	06-05-1994
		CN 1090581 A,B	10-08-1994
		DE 69301897 D	25-04-1996
		DK 596802 T	03-06-1996
		ES 2085130 T	16-05-1996
		GR 3019385 T	30-06-1996
		HU 66446 A	28-11-1994
		JP 6220082 A	09-08-1994
		US 5527780 A	18-06-1996
		US 5614614 A	25-03-1997
WO 9731929 A	04-09-1997	FR 2745290 A	29-08-1997
		AU 1929897 A	16-09-1997
		EP 0885234 A	23-12-1998